

**UNITED KINGDOM**

**RESEARCH ON GEODESY**

**1999 – 2002**

# **UNITED KINGDOM RESEARCH ON GEODESY 1999 – 2002**

**Prepared on behalf of  
The Royal Society's International Relations Committee**

**Submitted to  
The International Association of Geodesy  
at the XXIII General Assembly of IUGG  
Sapporo, Japan, July 2003**

**Published on behalf of  
The Royal Society  
by**



**The Institute of Engineering Surveying  
and Space Geodesy  
The University of Nottingham  
University Park, Nottingham, NG7 2RD**



## FOREWORD

This report outlines United Kingdom activities in geodesy for the period January 1999 to December 2002. It has been prepared for submission to the International Association of Geodesy (IAG) at its General Assembly in Sapporo, Japan, during the XXIII General Assembly of the International Union of Geodesy and Geophysics (IUGG) in July 2003. It is issued on behalf of the Royal Society.

Following the pattern set by the previous UK National Geodesy Report this document is presented in an undivided form, i.e. it is not structured to reflect the five sections of the IAG. The objective of this is to emphasize the linkage between the various disciplines within geodesy and to avoid earlier difficulties in assigning certain activities to particular sections. The document has been prepared within the Institute of Engineering Surveying and Space Geodesy at the University of Nottingham from information provided by UK geodesists.

There have been no significant changes in the structure of geodesy within the UK since the publication of the last report. The Geodesy group within the RAS/GS Joint Association for Geophysics has not however been effective as a means of communication amongst the rather small group of "geodesists" in the UK. Most communication and the dissemination of geodetic information has been facilitated through mail base user groups for "geodesy", "satellite navigation" and "geomatics", and through journals. There is not an effective national forum for geodetic matters, and this is a matter of some regret.

The editor wishes to thank Dr David Baker for his work in collating information and for preparing this report, and the Royal Society for agreeing to fund its publication.

ALAN H DODSON  
Editor and National Correspondent to the IAG  
June 2003

## CONTENTS

1. Satellite Laser Ranging	3
2. Global Navigation Satellite Systems	4
a. Atmospheric Studies	8
b. Engineering Applications	9
3. National and Continental Networks	12
4. Integrated Systems and Inertial Navigation Systems	14
5. Satellite Altimetry	15
6. Synthetic Aperture Radar	16
7. Satellite Orbit and Gravity Field Determination	17
8. Gravity Surveys	18
9. Theoretical Geodesy, Earth Tides, Earth Rotation and Miscellaneous Gravimetric Studies	18
10. Geoid Determination	20
11. Deformation Monitoring	22
12. Mean Sea Level Studies	23
13. Geophysical and Oceanographic Applications of GNSS	25
14. Law of the Sea	26
Bibliography	27

# 1 SATELLITE LASER RANGING

- **Department of Geomatic Engineering, University College London,**  
<http://www.ge.ucl.ac.uk/>

**SLR system calibration, Herstmonceux:** The SLR facility at Herstmonceux uses local calibration targets (LCTs) to determine system range errors for correction of its space observations. Current LCT ranges are based on a survey carried out by OS in 1991. Since the time of the survey the LCT positions have changed and range information has been updated by direct transfer using the laser ranger. The NERC Space Geodesy Facility at Herstmonceux required independent verification of these new calibration values and this was carried out by the Department of Geomatic Engineering at UCL. Leica System 500 GPS receivers were used over very long observation periods to establish a high-precision survey network relating the LCTs to two ITRF reference points and further reference marks in the vicinity of the SLR telescope (*Md Kadzim, 2002*). Theodolite intersection based on the above reference marks then measured the invariant point of the SLR telescope under rotation and thereby determined its position and distance from the local calibration targets (*Novak, 2002*). Accuracy is estimated to be better than 0.002m.

- **The NERC Space Geodesy Facility (NSGF),** <http://nercslr.nmt.ac.uk/>

**Satellite laser ranging:** The satellite laser ranging system of the NSGF at Herstmonceux has continued in regular operation throughout 1999-2002. It is operated, weather permitting, in a flexible manner, 24 hours a day, seven days a week. Given UK weather, SGF is recognised as one of the world's most efficient and productive SLR systems.

The system makes very accurate range measurements to Earth-orbiting satellites by timing the flight of short laser pulses from the telescope to the satellite and back. The target satellites carry retro-reflectors specifically for this purpose, and a range measurement is made about once every second during each satellite pass, with a precision of at best 8mm.

The UK SLR system belongs to a global network of stations coordinated by the International Laser Ranging Service (ILRS), one of 9 Services overseen by the International Association of Geodesy (IAG). Currently this global network tracks 22 satellites whose scientific missions fall into five broad categories:

- 1) Dedicated geodetic satellites (e.g. LAGEOS, ETALON) - for research into, and maintenance of scale and origin of, a global geocentric International Terrestrial Reference Frame (ITRF) at mm levels of precision;
- 2) Altimeter and Synthetic Aperture Radar (SAR) remote sensing EO satellites (e.g. ENVISAT, JASON-1, ERS-2) - tracked to determine accurate orbits and calibrate on-board sensors;
- 3) Dedicated gravity-field missions (e.g. CHAMP, GRACE) - use onboard GPS for precise orbit determination and SLR for orbit determination, stability and validation;
- 4) Navigation satellites from the GNSS constellations (e.g. 2 of GPS, several GLONASS) - tracked to give an independent measurement of the accuracy of their orbit determinations;
- 5) Experimental satellites - for testing new designs of reflector arrays, etc.

The ILRS sets performance targets, in terms of the precision of the measurements and numbers of passes tracked, for systems to be considered fully operational ILRS stations. Of the 40 stations registered with ILRS, some 15, including SGF, provide the bulk of the high-quality, high volume tracking for precise orbit determination necessary to meet the scientific goals of the satellite missions.

SGF, in its role as an ILRS Associate Analysis Centre, is collaborating with research groups from some 10 institutes towards an official, regular, ILRS station coordinate and EOP product. During 2002 a related 'Call for Participation' paper was jointly developed and distributed to the ILRS analysis community.

SGF continues to deliver daily, via its website, an automatic, orbit-based quality check on SLR data from the Herstmonceux system and from the global tracking network. This service enables rapid, independent feedback of SLR data quality and quantity for the benefit of both the SGF and the wider community.

In order to maintain the SLR system in a competitive position within the ILRS network, a level of research and development is carried out by the group and with international collaborators.

The SGF has made important discoveries about the characteristics of individual time-of-flight counters used to make the satellite range measurements. Similar careful experimentation and modelling has led to a greater understanding of the behaviour of the single photon avalanche diode (SPAD) detector, widely used elsewhere in the global network. Unless properly calibrated and operated strictly at single photon levels of return, standard practice at Herstmonceux, SPADs can induce significant bias into the measurements.

SGF and collaborators, within the ILRS Signal Processing Working Group, have made significant progress in the determination of precise values of the corrections required to transfer SLR ranges from the effective reflection points on satellites to their centres of mass, as functions of tracking system characteristics.

## 2 Global Navigation Satellite Systems

- **School of Civil Engineering and Geosciences, The University of Newcastle,**  
<http://www.ceg.ncl.ac.uk/>

**GNSS:** Investigations into sub-daily GPS coordinate solutions affected by un-modelled vertical signals revealed that biases are present in the horizontal coordinates when the ambiguities are not resolved as integer values (*King et al., to be published in 2003*). Biases are greatest in the east coordinate and are 40-50% of the magnitude of the vertical signal for semi-diurnal frequencies. A sensitivity analysis revealed that the ambiguity parameters were absorbing the un-modelled vertical signal and the horizontal expression of this is counter-balanced by biases in the horizontal coordinates. Resolving ambiguities or modelling the vertical signal eliminated the biases.

Sensitivity analyses were carried out on theoretical and proposed GNSS satellite constellations. Precision and reliability changes were investigated for a number of combinations of LEO, MEO and geostationary GNSS satellites, with relevance to the proposed Galileo system (*Rainbow, 2002; Rainbow & Clarke, 2002*). It is found that a combination of MEO and geostationary satellites offers significantly better precision and reliability than a purely-MEO constellation.

We have investigated the use of small GPS networks for determining tropospheric zenith delay and its consequent application to the correction of tropospheric artefacts in InSAR interferograms (*Wadge et al., 2002*). Our findings are that in certain cases the tropospheric information obtainable from small networks is of value to removing artefacts in interferograms, especially in mountainous regions where the tropospheric delay is highly correlated with surface elevation.

Low-cost GPS L1 phase receivers have potential for widespread use in vehicle tracking and attitude monitoring, but this field remains in its infancy. We have carried out trials of these receivers using single-epoch positioning and ambiguity resolution algorithms previously developed at the University of Newcastle. Initial results show that freely-floating solutions do not succeed due to poor a priori coordinates and high phase noise, but that solutions constrained with a priori information can have moderate success. We have also investigated the combination of GNSS with inertial and video data using Kalman filtering (*Carroll, 1999*).

- **Department of Geomatic Engineering, University College London,**  
<http://www.ge.ucl.ac.uk/>

**GPS phase multipath studies:** The Department of Geomatic Engineering at UCL has investigated the use of signal to noise ratios output by standard (Costas) GPS receiver phase tracking loops to model multipath errors in double differenced phase measurements. By using non-standard Fourier analysis techniques it was discovered, and reported in Cross, 1999, that in static environments around a 50% improvement in noise levels could be made through such modelling and, more importantly, that the long term error signals were virtually totally removed – resulting in errors with almost white noise characteristics. Methods of assessing multipath, and other aspects of GPS performance, were reported in Edwards et al, 1999. On-going work is concentrating on using output from more sophisticated phase tracking loops to estimate more directly the size of multipath errors.

- **IESSG, The University of Nottingham,** <http://www.nottingham.ac.uk/iessg/>

**POLARIS:** Polaris is a new EC-funded project to develop a software tool that will provide a link between the Galileo System design and users of the system. Polaris analyses the Navigation Performance of Galileo, and allows users to test the performance when augmented by various other sensors, such as odometers, compasses, and even GPS. Polaris includes a graphing utility with an interface to an internal GIS application.

Polaris will help identify new GNSS services, optimise the Galileo system from a user point of view, and optimise the Galileo Service Definition. Polaris will be a powerful means of illustrating the role of Galileo and showing users its potential benefits.

The Polaris team is led by GMV in Spain. The IESSG leads the work package on the User Application Subsystem, and is responsible for modelling the environment and trajectory of specific applications, and for providing detailed modelling algorithms for the additional sensors.

**GalileoSat:** In 2000, the IESSG completed a package of work as part of the ESA funded GalileoSat project led by Alenia in Italy, which studied the technical details of the space and ground segments of the future Galileo system. The IESSG's task, under the leadership of GMV in Spain, and coordinated by Alcatel Space Industries in France, was to study the definition, establishment and maintenance of the coordinate reference frame that would form the basis of all coordinates from Galileo.

**GALA:** Under a contract to Racal Tracs in the UK, and as part of the Alcatel-led team working on an EC contract, the IESSG studied the scientific market for Galileo services and receivers. The study was part of a much wider work package which investigated all potential markets for Galileo-enabled services, in order to extract user requirements and to estimate the potential revenues from Galileo services. It involved a great many contractors from across Europe, each tasked with studying the market for a particular segment of the market.

**The GalileoSat System Simulation Facility project (GSSF):** GSSF is a contract funded by the European Space Agency (ESA) to produce an end-to-end software simulation of Europe's planned 'Galileo' satellite navigation system. It will model all of the components of the future system, from the satellites themselves, to the ground segment that will control them and the user receivers that will navigate with their signals.

As part of the VEGA-led team working on the study, the University of Nottingham is acting as a consultant on technical aspects of satellite navigation and specific modelling details. Nottingham's role is to provide definitions of suitable simulation and modelling algorithms, with the necessary algorithm description, pseudo-code where necessary, and test data. One of the future aims of GSSF is to be able to replace software components of the system with real hardware - a so-called hardware-in-the-loop design. For instance, early versions of GSSF will simulate a Galileo user receiver in software, but later versions may wish to replace the software receiver with a prototype hardware version. This requires that the simulation models are as true to life as can practicably be achieved.

Following delivery of the phase one version of the software, the project is making good progress towards a version with high fidelity models of all aspects of the system. It will allow system engineers to investigate a wide range of Galileo issues, from the global performance of the system, to the effect of a small change in a particular hardware component.

**The British Isles GPS archive Facility (BIGF)**, <http://www.bigf.ac.uk>: BIGF, operating since 1998, is the long term repository for continuous GPS data recorded by a network of over fifty permanent GPS stations (including the Ordnance Survey active stations), established throughout the British Isles. It has recently gained funding from the Natural Environment Research Council (NERC) to become one of their prestigious facilities. BIGF is hosted by the IESSG, known globally as a centre for postgraduate teaching and research in the field of satellite positioning. It therefore provides an appropriate venue for a central archive of such data.

By archiving GPS data on a long term basis, BIGF preserves long term signatures buried in the data, also enabling cost economics when bidding for research funding.

BIGF facilitates research by non-commercial scientific users for use in scientific applications, recent examples include:

- *Kinematic data processing*
- *Precise height determination*
- *Atmospheric water vapour studies*
- *Ionospheric studies*
- *Numerical weather prediction*
- *Signal delays due to the atmosphere*

The aim of BIGF is to *enable access* and to *broaden the range* of disciplines using the archive. Access to the archive is enabled firstly by submitting an online form at the BIGF website, with data delivery by secure ftp.

- **Imperial College London**, <http://www.geomatics.cv.ic.ac.uk>, <http://www.cts.cv.ic.ac.uk>

**Performance potential of a combined Galileo/GPS navigation system: This project was carried out for Alcatel Space by the London Centre for GNSS Research (LCGR):** The LCGR is a joint research initiative by Imperial College London and University College London. The objectives of the research project were to define an integrity monitoring concept for the Global Positioning System (GPS) satellites within the Galileo system, and to quantify the level of *improvement (accuracy and integrity)* to be gained by the adoption of a combined Galileo/GPS navigation system. This was part of a wider study by Alcatel Space aimed at the definition of the Galileo system. The results were used as input to the design of the Galileo system and in international negotiations with the United States of America and other countries.

**EGNOS expansion:** This project was carried out for the European Space Agency (ESA) to study the requirements for the expansion of the European satellite navigation system, EGNOS (European Geostationary Navigation Overlay Service) to Africa and South America. Imperial College London was a member of a multinational consortium with the responsibility for developing the algorithms and code for modelling navigation system errors required for the expansion studies.



- **The NERC Space Geodesy Facility (NSGF)**, <http://nercslr.nmt.ac.uk/>

**GNSS:** SGF manages two IGS systems at Herstmonceux; HERS, currently an Ashtech Z12 system, operating since 1996 and HERP, a joint GPS/GLONASS dual frequency Z18 receiver.

During 2001 an antenna fault was discovered on the HERS system and since rectification the data quality and consistency has improved dramatically. From 2001, the GNSS observations are rigorously checked on site for consistency and quality and then delivered in accordance with the IGS guidelines; the GPS and GPS/GLONASS data both hourly and daily are delivered in standard RINEX format to IGS data centres.

In order to improve the low-elevation sky coverage seen by the GPS/GLONASS system HERP, the antenna will be moved during 2003 from its current location to a superior location close to the principal ground-calibration target used by the SLR system. The move will further strengthen the local links between the GNSS and SLR co-ordinate frames. As part of an ongoing programme of local surveys (a collaboration between SGF and the Geomatics Department at UCL) two MSc students carried out a survey during the summer of 2002. It is likely that the re-sited HERP system will be also be used in support of LEO satellite missions and to monitor the Earth's atmosphere in near real-time, via 1 Hz broadcasting through the Internet.

Evaluation of potential systematic bias in GNSS orbital solutions (SGF): The high accuracy of SLR measurements can be exploited to test the accuracy of operational orbital ephemerides of GNSS satellites, as determined by IGS. Two GPS satellites and all GLONASS satellites are fitted with laser retro-reflector arrays and the global network makes precise ranging observations on a regular basis. Because the GLONASS satellites carry large arrays the interpretation of SLR observations is not entirely straightforward. But detailed research has shown that, with an uncertainty of about 2cm, the laser range observations are a good test of the quality of the radial component of the IGS-computed orbits. For the small arrays on GPS satellites there is no such problem of interpretation. The laser data are consistent in showing that the IGS orbits are too large; ranges determined from the IGS orbits place the satellites about 5cm further away from the centre of the Earth than the SLR observations imply. The radial scatter in GLONASS orbits is found to be about 20cm RMS and that of the GPS orbits is about 12cm RMS.

- **Proudman Oceanographic Laboratory**, <http://www.pol.ac.uk>

**GPS time series analysis:** Until recently, it was typically assumed that only white noise was present in geodetic time series. However, several data sets of continuous GPS time series have now provided evidence for the presence of time-correlated (coloured) noise. We have shown that the presence of coloured noise can significantly affect the uncertainties of rates estimated from these time series. We derived a set of empirical equations to account for the coloured noise. Our analysis of time series from GPS sites around the world and regional networks have further confirmed the presence of coloured noise in these data sets and that there is a latitude dependence (and hemispheric bias), a time dependence (gradually decreasing) and a spatial correlation in the noise. Regional networks can take advantage of the spatial correlation to reduce the amount of noise present in each time series. Investigation into the role offsets play in the estimation of site velocities has been conducted. Analysis of present GPS data sets reveals that, on average, one offset occurs every ten years, but it may be as often as once every two years. We have derived strategies and equations for dealing with offsets that will both limit the effect they have on site velocities and account for them in the velocity uncertainties.

## 2a Atmospheric Studies

- IESSG, The University of Nottingham, <http://www.nottingham.ac.uk/iessg/>

**Ionospheric scintillation effects on GPS measurements:** The ionosphere is prone to significant disturbances, which are considerably worse during periods of high solar activity, such as the recent solar maximum. Ionospheric scintillation, the most significant manifestation of such disturbances, often takes place in equatorial, and auroral regions such as Northern Europe. Since June 2001 a network of state-of-the-art GPS Ionospheric Scintillation Monitor (the NovAtel/AJ Systems GSV4004) receivers is being used to measure and record GPS phase and amplitude scintillation parameters observed in the UK and in Norway. These specially designed GPS receivers have been co-located with permanently tracking dual-frequency GPS receivers, forming a Northern European monitoring network. This is the core of a 3 year EPSRC (Engineering and Physical Sciences Research Council) funded project, running at the IESSG. The aim is to study the effects of ionospheric scintillation and Total Electron Content (TEC) gradients on GPS applications during the solar maximum, with a focus on accuracy, availability and integrity.

The project stages involve the set up of the network, the development of automated archiving and data analysis strategies, impact assessment of ionospheric scintillation on maritime DGPS and EGNOS users, and on different GPS receiver technologies. In addition to the monitoring network, data obtained from existing International GPS Service (IGS) stations and from the General Lighthouse Authorities (GLA) DGPS reference stations have also been used. Analyses have been carried out on data from periods of known ionospheric activity, but long term systematic studies will also be undertaken. The GPS user errors have been correlated with observed scintillation levels and with available geomagnetic indices during various active periods. Results of the impact assessment exercise, both on the DGPS and EGNOS user, have been published. The study of the impact on different GPS receiver technologies is to be carried out in 2003. On the background of the main objectives of the project resides the idea of forming a long term data base for use by the scientific community and the possible development of warning and/or mitigation mechanisms for the GNSS user community. We have found that amplitude and phase scintillation levels predicted by the WBMOD (Wide Band Model) ionospheric scintillation prediction program present good correlation with GPS user errors and the possibility of mitigating the GPS user errors with the aid of this model is currently under investigation. The scintillation monitor receiver outputs raw scintillation data at a rate of 50Hz, but is also capable of recording only the relevant statistics of these data at every 60 seconds, at different intervals. As a baseline these statistics have been recorded and used in our analysis. However a high rate campaign, with the recording of the 50Hz raw data, is planned for 2003. Please see <http://www.nottingham.ac.uk/iessg/isgres38.htm>

**Meteorology and climate:** The IESSG continued its research into the use of GNSS signals for meteorology and climate studies. In particular a collaborative project with the UK Meteorological Office (within the general context of the EC COST Action 716) was initiated, aiming to provide a UK operational capability. Research was particularly focussed on the ability of GNSS network to monitor severe weather events, such as thunderstorms. In addition the Institute continued its research into atmospheric delay estimation from receivers on moving platforms. A collaborative project with the NERC ESSC in Reading investigated GPS delay estimation above and around Mount Etna, Italy, in an attempt to remove atmospheric biases from Interferometric SAR observations.

**River level monitoring:** The RiGHt project "River Level Monitoring using GPS Heighting" was completed in 2002 by a consortium comprising Science System (Space) Ltd, the Institute of Engineering Surveying and Space Geodesy of the University of Nottingham and the Centre for Ecology and Hydrology. The two phases of the project were both funded by the British National Space Centre (BNSC), under the UK Government's Space Foresight Programme.

The original RiGHt project, which ran for two years, was successful in demonstrating the feasibility of measuring and monitoring river heights using a GPS equipped buoy in a real-time environment and using satellite communications technology to transfer the data to a central and remote Geographical Information System.

The aim of RiGHt has been to demonstrate innovation in the integration of state-of-the-art technologies such as OTF GPS and low power world-wide satellite communications to solve a real user need. The objectives of the second phase were to address three key developments to the original architecture. Firstly, the use of long-range kinematic GPS techniques was implemented, so as to extend the useful range of the system. Secondly, long-range use requires the adoption of satellite communication techniques to relay the raw data between the buoy, other reference GPS stations and a central computational hub. This has required the use of advanced data compression techniques, to handle the high data rates necessary for raw GPS carrier phase data. The third objective was to integrate the resulting real-time measures of river level into river and flood modelling software, so as to greatly enhance the benefit of the monitoring to the users.

The development of the new RiGHt system was completed by early 2002, and in March a series of land and river based trials were conducted. The RiGHt project has been highly successful. The system may now be deployed in routine monitoring applications or in situations of real emergency. It can provide a very elegant and rapid solution to many users' needs.

## 2b Engineering Applications

- **Department of Geomatic Engineering, University College London,**  
<http://www.ge.ucl.ac.uk/>

**GPS attitude determination:** Several GPS attitude studies have been undertaken in the Department of Geomatic Engineering at UCL. One involved integration of several sensors, including a multi-antenna and multi-receiver GPS-based system, to provide both attitude and position for airborne remote sensing operations. It was applied to data collected on board the UK Natural Environment Research Council's Airborne Remote Sensing Facility aircraft (Sheridan et al 2002, and Sheridan and Cross, 2000). Another project involved the design and testing of an attitude determination algorithm for a low orbiting spacecraft using GPS signals as the sole observable (*Jameson et al, 2001, and Cross and Ziebart, 2002*). The algorithm was designed to run in real-time at a rate of 10Hz on-board the spacecraft, using only minimal chip and memory resources. Principal features of the algorithm design were: rearrangement of the conventional double difference GPS phase observation equations in terms of the Euler angles alone (thus reducing the number of solve-for parameters to three), numerical differentiation of the model equations, no requirement for a Kalman Filter, a novel single epoch ambiguity resolution technique and the facility to trap and fix cycle slips. Testing was effected by developing an independent observation simulator.

- **IESSG, The University of Nottingham,** <http://www.nottingham.ac.uk/iessg/>

**Combined use of virtual scale models and space geodesy:** Late in 2001 the Institute was awarded a ROPA (Rewarding Our Potential) grant from the Engineering and Physical Sciences Research Council to develop blue sky research on an area that exploits the existing expertise of the IESSG in high precision GPS positioning and an equally specialised know-how, existing at the University's MRL (Mixed Reality Laboratory), in augmented reality. The aim is to investigate and develop a means to give engineers, and the community affected by civil engineering work, information that is often difficult to visualize via computer screens or 2D plans. At the end of 2002 a prototype system had been developed, in which the user works in a collaborative environment where it is possible to see virtual and real features of a test-bed area seamlessly in real time. With this information, the users of the system can assess and discuss how the environment will be affected by the engineering work, enabling a vision of the future and the possible solution of problems that could appear during construction, further enabling the pursuit of cost reduction and improvements in efficiency.

**Augmented reality for sub-surface visualisation:** The IESSG in collaboration with the School of Chemical, Environmental and Mining Engineering were sponsored by the Department of Trade and Industry to develop a GPS based Augmented Reality system. Ten external organisations were also involved in the research. Augmented Reality is the science of superimposing Virtual

Reality data onto the real world, in real time. Applications developed during field trials include superposition of buried pipe and cable information, mining and geology information onto real world views.

**Bridge deformation monitoring with GPS:** The overall of the project is the creation of a system that employs computational simulation with GPS to monitor the health of bridges without the need for onsite inspection. The main research interest is GPS and making it more effective for the specific aim of bridge deflection monitoring. GPS has been proved in the past to be an efficient tool for this monitoring, but research is still being conducted into mitigating the associated error sources and providing more accurate deformation information. In the past expensive dual frequency receivers have been used with good results, but one of the aims of this project is to use cheaper single frequency receivers. The aim is the creation of a more affordable monitoring system. Displacements that deviate from the normal design configurations will redistribute stresses and strains amongst the other bridge components, which will in turn affect the load carrying capacity of the whole bridge. A system that can spot abnormal movement or changes in frequency of the bridge can be used to flag faults that could lead to catastrophic failure.

Since starting research the author's main interests have been accelerating the ambiguity resolution in single frequency receivers, so that cheaper receivers can be used for this application; using a total station for dynamic bridge monitoring and comparing the results to GPS; and introducing pseudolites (pseudo-satellites) to improve the reliability, integrity and accuracy of the solution. A number of bridge trials have taken place, the results of these have been processed and analysed.

**Detection of abandoned mineshafts:** This is a collaborative research project with British Geological Survey supported by the EPSRC, involving the development of a vehicle-towed capacitively coupled form of resistivity detector, to be used in searching for abandoned mineshafts. Unlike many geophysical sensors this system can take measurements whilst moving, enabling high rates of data capture, but requiring highly accurate positioning, which can only be provided by RTK GPS. To minimise costs a single frequency GPS receiver can be used.

The Institute has now developed the real-time navigation, triggering and positioning element of the geophysical equipment, which has been field tested on buried objects at the BGS and over known mineshafts at Bonstal moor.

**Construction plant control and guidance using RTK GPS:** Work in this area continues through a project part funded by Trimble Navigation Europe Ltd. The research work includes a full scale trial where a GPS system guided and controlled a bulldozer whilst clearing and levelling an area of ground. The project aims to assess the strengths and weaknesses of using GPS in plant guidance and examines its current and future roles in the construction industry. Testing was carried using a Trimble provided SiteVision GPS plant guidance/control system, including quality assessment of individual system components and assessment of performance as a whole in a full scale trial.

GPS signals used in plant control applications, in precision grading for example that requires carrier phase measurements, suffer from multiple error sources, in particular signal multipath, causing measurement error of up to 5 cm. Whilst many techniques exist to mitigate pseudorange measurement multipath, or carrier phase multipath in static applications, the issue of carrier phase multipath for a dynamic receiver has not been adequately investigated. Part of this project therefore, involves research into the efficiency of various methods of identifying and mitigating carrier phase multipath in real-time for dynamic applications, and furthermore how GPS can be augmented in difficult signal reception environments using inertial sensors, and other positioning systems.

- **Imperial College London**, <http://www.geomatics.cv.ic.ac.uk>, <http://www.cts.cv.ic.ac.uk>

**Transport Telematics - Local area augmentation system for fleet management:** This project was funded by Logica (UK) Limited to specify an architecture for a local area navigation system for fleet management in urban areas. The architecture proposed consisted of the use of Medium Frequency (MF) signals generated from terrestrial MF signal generators (similar to those used for LORAN-C) to augment the signals from the space based navigation systems such as GPS, GLONASS and Galileo.

**Transport Telematics - Space-based dynamic positioning in built-up environments:** This is an ongoing project on high accuracy and high integrity dynamic positioning in urban areas to support road transport navigation. Part of this has been the contribution of Imperial College London to the review of the Countdown Bus Information System operated by Transport for London (TfL). The project assessed the capability of stand-alone and augmented space-based navigation systems to support the navigation functionality of the Countdown system. Currently, the system uses roadside beacons and distance information from odometer to determine the location of buses. Research is continuing to investigate data fusion techniques involving the integration of GNSS with low-cost MEMS technology sensors to enable required navigation performance in built-up areas.

**Transport Telematics - The development and demonstration of a vehicle performance and emissions monitoring system:** This is a *Foresight Vehicle LINK* project awarded to a consortium including Imperial College London, Sira Technologies Ltd and Saturn Technologies Ltd, to develop a real-time vehicle performance and emission monitoring system (VPEMS). It will be completed in 2003 with funds from the UK government and industry. The project is developing further and applying state-of-the art technologies to create and demonstrate the capabilities of an accurate, reliable and cost effective VPEMS. The system is to be fitted on vehicles to monitor driver/vehicle performance and the level of concentration and emissions inside the vehicle and at the exhaust respectively. Real-time sensing of spatially and temporally referenced vehicle performance and emission data will enable, for example, the monitoring of environmental compliance, identification of polluters and effective management of maintenance of vehicles. Such data will also facilitate research on the major drivers of vehicular pollution and the corresponding health impacts. A special advisory committee consisting of colleagues from industry and government agencies is facilitating the development and exploitation activities.

**Transport Telematics - Measures of road traffic congestion:** This project is exploring the extent to which existing data sources including GNSS, could be used to improve on the methods for measuring traffic congestion. The project is divided into two phases, one to survey existing data sources and the other to investigate how the data could be used to provide a reliable and relevant measure or measures of traffic congestion. Issues of coverage (spatial and temporal) and data quality are important particularly if data from different sources is to be required.

**GPS integrity and air traffic service hazard assessment:** The objective of the first phase of this project was to study the strengths and weaknesses of the global positioning system (GPS) of satellites as a navigation tool for civil aircraft. The study quantified the level of integrity (safety) afforded by GPS both at system and user levels, related this to civil aircraft navigation requirements and finally, proposed techniques for improving the performance of GPS to meet the requirements. The next phase of the project will carry out a hazard assessment of GNSS-based air traffic services.

**Integrated high precision kinematic positioning using GPS and EGNOS observations:** This project has developed algorithms and software to assess the performance achievable when combining range and carrier phase measurements from different satellite navigation systems, GPS and Geostationary Earth Orbiting (GEO) satellite data within the EGNOS system. Additional satellite systems are required to augment GPS in order to increase the performance (accuracy, integrity and availability). The results showed a major improvement in real-time kinematic positioning when GPS data is combined with EGNOS data.

**Structural stability monitoring using satellite positioning systems and GIS:** The objective of this project is to specify and demonstrate a system architecture for structural deformation

monitoring based on satellite positioning and geographical information systems technologies, and to develop methods for using this data to understand structural behaviour. The work requires a multi-disciplinary approach combining advanced satellite positioning and geographical information systems technologies with background in structural analysis and assessment as well as external input from practicing engineering firms.

**Earthquake risk assessment using remote sensing:** A consortium consisting of Imperial College London and European industry was awarded a contract by the European Space Agency (ESA) under the Long Term Development of Earth Observation Market initiative. The project is developing and validating the technique of Permanent Scatterer Interferometry Synthetic Aperture Radar (PSInSAR) for the purpose of deriving high precision maps of ground displacements over time. Imperial College London is contributing expertise in positioning with GPS to validate the new PSInSAR concept.

**Air Traffic Control and Management - Assessment of the impact of future air traffic management technologies and procedures on airspace capacity:** With global demand for air transport doubling every 10-15 years, the total number of flights in Western Europe is forecast double between 2000 and 2010. Current air traffic management (ATM) procedures and technology cannot cope. There are increasing safety risks, with growing numbers of recorded 'near misses', and ever increasing costs of delay due to congestion. The technological challenge is to increase airspace capacity without compromising safety, environmental and economic requirements. An on-going project supported by the EPSRC and the Royal Society is exploring the impact of new technologies (in particular satellite navigation systems) and management procedures on airspace capacity. The involvement of the ATM industry is crucial to the success of this project. Eurocontrol and QinetiQ are collaborators on the project.

**Air Traffic Control and Management - Cross-sectional time-series analysis using simulated controller workload data:** This project is funded by Eurocontrol to carry out further research on understanding airspace capacity's dependence on controller workload. A cross-sectional time-series analysis using simulated controller workload data will be carried out.

### 3 NATIONAL AND CONTINENTAL NETWORKS

- **IESSG, The University of Nottingham**, <http://www.nottingham.ac.uk/iessg/>  
**Ordnance Survey of Northern Ireland**, <http://www.osni.gov.uk/>  
**Ordnance Survey Ireland**, <http://www.osi.ie/>  
**Commissioners of Irish Lights**, <http://www.cil.ie>

**IRENET:** The active GPS network for Ireland is the result of collaboration between Ordnance Survey Ireland (OSi), the Commissioners of Irish Lights (CIL), and the Ordnance Survey of Northern Ireland (OSNI) and comprises of sixteen stations distributed around the island. During the first half of 2002 an extensive GPS measurement campaign was undertaken to connect the sixteen GPS network stations to the European GPS co-ordinate reference framework (EUREF). The IESSG processed the GPS data to determine accurate ETRS89 coordinates for the 16 active GPS stations, which will become the realisation of ETRS89 in Ireland. In addition to this, assessments of the station velocities and movements of the 11 IRENET95 Zero Order passive stations and the 3 EUVN97 passive stations were made. The ETRS89 coordinates of a network of 29 FMB and tide gauge passive station were also determined, relative to the active network stations.

- **International Institution for History of Surveying & Measurement (A body within FIG).**

**Struve geodetic arc and its extension:** Since 1994 the History of Surveying Group within FIG, with the support of IAG and IAU, has been working on a project to get a selection of the remaining surveys stations on the Arc recognised as an UNESCO World Heritage Monument.

The Arc was observed between 1816 and 1855 and contained 265 station points. In today's geography it passes through ten countries :- Norway, Sweden, Finland, Russia, Estonia, Latvia, Lithuania, Belarus, Ukraine and Moldova.

It has been a considerable challenge to get ten countries to each produce all the necessary field work and documentation but all is gradually coming together.

The aim is to finish with between 35 and 40 points preserved and suitably marked. Submissions to UNESCO can only be made once a year and it is hoped to submit the case for the Arc in February 2004. The National Land Survey of Finland has kindly agreed to collect, collate and print all the necessary documentation.

Following on from this project efforts are now being directed to verifying a link from the southern end of the Struve Arc (Izmail on the Black Sea) via Romania, Bulgaria and Greece to Crete then across the Mediterranean to Egypt and thence through the Arc of 30th Meridian to South Africa.

If successful, it is hoped to apply a similar World Heritage Monument exercise to this triangulation and thus form a monument from North Cape to Cape Aghulas - a distance of some 11 450 km or almost 105 degrees.

If any readers are aware of information relating to a connection from Izmail to Crete. either circa 1868 or later, we would be very interested to hear about it. Please contact J R Smith, Honorary Secretary to the Institution, at [101765.332@compuserve.com](mailto:101765.332@compuserve.com)

- o **Ordnance Survey of Great Britain**, <http://www.ordsvy.gov.uk>

**New official extension to the EUREF for Great Britain, EUREF GB 2001:** The purpose of the EUREF GB 2001 campaign was to compute new ETRS89 coordinates for all the stations in the Ordnance Survey Active GPS Network. Although GB had an existing network of EUREF stations (EUREF GB92), this was based upon ground marked stations only and did not include any Continuously Operating GPS Receivers (COGRs). The EUREF GB 2001 campaign was based upon two weeks of observations at the 30 permanent GPS stations which comprise the Active GPS Network, four additional stations and six fiducial IGS stations in Europe. The coordinates were computed following the latest recommended methods of the EUREF Technical Working Group. 20 stations from the campaign were proposed and officially ratified at the EUREF 2002 symposium as an improvement and extension of EUREF89.

**Ordnance Survey National GPS network**, <http://www.gps.gov.uk>: Launched in 2000, the website gives online users free access to: the Ordnance Survey Active GPS Network RINEX data server; the passive GPS station database; online coordinate converter (utilising the OSTN02 / OSGM02 transformations) and information resource.

The Active GPS network consists of 30 permanent geodetic GPS receivers installed throughout Great Britain, such that most locations are within 75 km of at least one Ordnance Survey active station, and major urban areas are served by several. All Ordnance Survey active stations record dual-frequency GPS data 24 hours a day at a 15 second epoch rate. The network control centre at Ordnance Survey headquarters in Southampton receives a one hour packet of GPS data from each active station once per hour around the clock, and adds this to the RINEX data server for immediate access by users.

The 900 passive GPS network station coordinates were updated following a complete readjustment of the entire network known as OSGPS2002. OSGPS2002 incorporates many extra observations and also extends the passive network into the Scottish Islands, the Scilly Isles and the Isle of Man. The expected coordinate errors at a passive station are now improved to 0.055 m horizontal (95%) and 0.066 m vertical (95%). Hundreds of observations between passive and active stations have been included in OSGPS2002 meaning that the two networks are now in closer sympathy.

**Definitive national grid transformation OSTN02:** OSTN02 is a transformation model which not only models the relationship between ETRS89 and OSGB36 but actually redefines OSGB36 in

terms of ETRS89. Since the introduction of the Ordnance Survey National GPS Network and the ratification of the Active stations as the official realisation of ETRS89 in Great Britain (EUREF GB 2001), the national geodetic coordinate frame has been defined in ETRS89. Prior to this, OSGB36 National Grid was realised by the coordinates of the old terrestrial Retriangulation. Because OSGB36 only exists on land, OSTN02 has been limited to within 10km offshore. OSTN02 is a grid look up type model and achieves an accuracy of 0.1m rms when comparing coordinates of lower order (third and fourth order) control achieved through GPS plus OSTN02 and existing archive OSGB36 coordinates.

- **United Kingdom Hydrographic Office (UKHO)**, <http://www.ukho.gov.uk>

**Control networks:** Work has continued to establish more geodetic control in the Antarctic Region areas of South Georgia, South Sandwich Islands, Antarctic Peninsula, South Orkney and South Shetland Islands as well as the Atlantic Island of Tristan da Cunha. Most data has been post-processed with IGS stations resulting in ITRF coordinates.

## 4 INTEGRATED SYSTEMS AND INERTIAL NAVIGATION SYSTEMS

- **IESSG, The University of Nottingham**, <http://www.nottingham.ac.uk/iessg/>

**GPS integrated with remote sensing:** GPS is being used in combination with terrestrial laser scanning and ground based photogrammetry for monitoring cliff erosion and slumping.

**Integration of GPS / INS:** The IESSG at the University of Nottingham have developed a sophisticated software package which enables the close and effective integration of RTK GPS with Inertial Measurement Units (IMU). An Adaptive Kalman Filter is used to jointly process the pseudo-range and carrier phase GPS measurements with the accelerometer and gyro readings. The IMU data greatly enhances the ability of the software to detect and repair cycle slips in the phase data and also greatly accelerate the ambiguity resolution. In a complementary manner the GPS data enables a far more rapid alignment of the IMU sensors and the containment of the biases and drifts of the sensors. The package has been tested with data from the IESSG Navigation Simulator and also real data from marine trials of the integrated system. Further work is now continuing to develop the software so that it may be exploited throughout a range of applications areas.

**Low-Cost Navigator (LCN):** The University of Nottingham is part of a team led by Thales working on the integration of GNSS and inertial sensors into a low cost package that is able to provide precise navigation in difficult environments. The University's role is to assist the design process by testing future components in a simulation environment. To achieve this, the University has developed a capability to simulate inertial components (gyros, accelerometers) as part of its GNSS simulator, in an integrated environment. Detailed models of the performance of the inertial components have been developed, so that synthetic GNSS data and inertial data can be produced from a common platform. A trajectory generator allows synthetic data to be produced from a combined GNSS/Inertial platform which is moving in a precisely controlled way. Data from this tool has been validated using an analysis filter from another consortium member, as well as with in-house analysis tools.



## 5 SATELLITE ALTIMETRY

- **School of Civil Engineering and Geosciences, The University of Newcastle,**  
<http://www.ceg.ncl.ac.uk/>

**Satellite altimetry:** We have continued with ongoing studies into the stability of altimetric range measurements using absolute and relative calibration techniques. An absolute calibration of TOPEX/Poseidon (*Dong et al, 2002*) and of ERS-2 (*Dong et al, to be published in 2003*) altimeter biases have been undertaken using UK tide gauges equipped with GPS. The tide gauge reading is extrapolated to sub-satellite points using a local geoid while the ocean tide numerical model has been replaced by estimation of the tide-difference parameters from the TOPEX data itself. The methodology is general and can be applied to other locations and is currently being used for Jason-1.

Relative calibration studies have been undertaken for TOPEX/Poseidon using the global network of tide gauges and for ERS-2 using dual crossovers with TOPEX/Poseidon (*Moore, 2001*). Intra-calibrations of TOPEX/Poseidon have revealed differences between Side A and Side B of the altimeter and a clear annual signature. A time series of the ERS-2 bias, as derived from dual crossover data, shows evidence of a small drift in the altimetric range with the sea-state bias a potential source of error.

- **Earth and Planetary Remote Sensing Laboratory, De Montfort University,**  
[http://www.dmu.ac.uk/faculties/cse/computing/eprsl\\_research\\_group.jsp](http://www.dmu.ac.uk/faculties/cse/computing/eprsl_research_group.jsp)

Over the past four years, research in the E.A.P.R.S. laboratory at De Montfort University has concentrated on the analysis, interpretation, and development of novel applications of satellite radar altimeter data. By interpreting the complex echo shapes obtained over non-ocean surfaces, a global database of over 1000 million orthometric heights has been generated from the ERS-1/ERS-2 altimeters, with additional data from Topex/Poseidon and., recently, Envisat RA-2.

One focus of research has been the investigation of existing Global Digital Elevation Models, both to determine the extent of errors by comparison with a globally distributed, independently derived height dataset, and to provide information on the effects of these errors on current global and regional gravity field models. Following on from this research, which confirmed the presence of very significant, regional scale height and terrain errors in current GDEMs, a new GDEM – ACE (Altimeter Corrected Elevations) was generated, by merging the best available ground truth with altimeter derived heights. Over 40% of the earth's land heights have been significantly altered in this radically different GDEM, which is now increasingly used, in applications as diverse as DTM 2002, and as the AATSR GDEM for Envisat.

In allied work, the potential for analysis and correction of regional DEMs was pursued, over Australia, in conjunction with Curtin University, Perth, and for SAR generated models, with the European Space Agency. Currently, the SRTM dataset is being assessed as the model is progressively released into the public domain: initial results over North America appear promising.

A new focus over the past four years has been the analysis of data over inland water. Initially working over river systems, the potential of retracked altimeter data to retrieve global information over inland water has been demonstrated.. Detailed river height data have been retrieved, and decadal time series have been analysed. Recently, this work has been extended to an analysis of global lake systems, and it has been shown that valid height data can be extracted from the complex altimeter echoes from ERS-1/2 for over three quarters of the earth's major lakes, and for one quarter of these lakes by retracking Topex data, allowing the derivation of global lake climatology. These time series are now being and analysed for climate-related information. With the ability of the new altimeter missions, Envisat and Jason-1, to release processed data quickly, this work is now being extended to near-real-time monitoring of inland water. Recognising the potential of this technique, a new high level hydrology product is being designed in collaboration with the European Space Agency. This will be available from 2004 for Envisat data.

**Envisat CAL/VAL:** One key activity during this period has been the involvement in the Calibration/Validation Team for the Envisat RA-2, both in investigation of the performance over land surfaces, and in analysis of the backscatter over natural land targets. Cross-calibration has been carried out with ERS-1 and ERS-2 ice and ocean mode backscatter, and Topex data are now being studied. The outcome of the Envisat Cal/Val activity shows that the RA-2 is retrieving data even over the extreme terrain of the Andes and Himalayas mountains. Of particular interest is the retrieval of the first ever altimeter heights from lake Titicaca, high in the Andes, demonstrating the potential for height generation and for monitoring of inland water even in extreme terrain.

**SSG 3.184 Use of remote sensing for monitoring heights and depths:** The E.A.P.R.S. lab has played a key role in the work of this IAG Special Study Group (chaired by Professor Berry) over the past four years, with a range of collaborative research projects with group members, and with development of new datasets such as the ACE GDEM.

- **Proudman Oceanographic Laboratory**, <http://www.pol.ac.uk>

**Satellite altimetry:** POL has continued to employ data sets from TOPEX/POSEIDON and ERS-1/2, and now JASON-1. Projects have included the use of altimeter data for ocean circulation studies, and research into the use of UK tide gauge equipped with GPS to provide an ongoing altimeter calibration system. A particular topic is the interaction of mean flows with eddies and Rossby waves. In the Southern Ocean it has been shown that the mean eastward flow overcomes the intrinsic westward propagation of Rossby waves to produce the world's only large region of eastward propagation of sea level anomalies, apart from the equator. The eddies in the Southern Ocean have been shown to remove eastward momentum from, or to produce a torque on, eastward jets. This is unlike the situation in many ocean models, suggesting a greater role for interactions with bottom topography than hitherto accounted for. The interaction between mean flows and transients has also permitted verification of the existence of zonal jets in the Coral Sea, predicted from model analyses. A comparison of seasonal cycles in the northern North Atlantic, as measured by altimetry and surface drifters, has also been undertaken, showing that the two datasets are consistent and complementary, and elucidating the flow pattern in this region which is important for the global thermohaline circulation.

Altimetry has also been compared with Antarctic tide gauge measurements (see the POL entry under *Mean Sea Level studies*).

## 6 SYNTHETIC APERTURE RADAR

- **IESSG, The University of Nottingham**, <http://www.nottingham.ac.uk/iessg/>

**Phase ambiguity determination:** A technique has been developed that resolves SAR phase integer ambiguity from single InSAR pairs. The method requires only very coarse, sparse ground control and has major implications for differential techniques and the derivation of atmospheric delay residuals.

## 7 SATELLITE ORBIT AND GRAVITY FIELD DETERMINATION

- **School of Civil Engineering and Geosciences, The University of Newcastle,**  
<http://www.ceg.ncl.ac.uk/>

**Satellite orbit and gravity field determination:** Precise orbit determinations have been undertaken for the altimetric satellites ERS-2 and TOPEX/Poseidon and their follow-on missions ENVISAT and Jason-1. Orbital analysis has also been used to develop a tailored gravity field for the ERS orbit (Moore, 2001) while dual and single satellite crossovers provide a measure of both the orbital accuracy and a mechanism for investigation of the relative altimetric bias. Work in this area has contributed to ESA's Orbit Validation Team (OVT) and Cross Calibration Validation Team for ENVISAT. Our in-house software FAUST has been extended to permit a reduced-dynamic solution which has proved superior to the standard dynamic methodology for ENVISAT.

Effort has been devoted to orbital analysis and gravity field recovery from the CHAMP mission. GPS tracking has been used within a reduced dynamic formulation of the GIPSY/OASIS II software to produce positioning for study of the gravity field and for participation in the IGS LEO Pilot Project (Moore et al, to be published in 2003). Errors affecting gravity field recovery from CHAMP have been investigated (Moore et al, to be published in 2003) with the conclusion that orbital positioning is the critical source of error with accelerometry contributing at a lower level. Attitude error is not significant.

- **Department of Geomatic Engineering, University College London,**  
<http://www.ge.ucl.ac.uk/>

**Analytical non-conservative force modelling for near-Earth satellites:** There is an on-going study in the Department of Geomatic Engineering at UCL to develop generalised methods of modelling the solar non-conservative force field (including effects due to solar radiation pressure, anisotropic thermal re-radiation and Earthshine/thermal emission effects). The techniques that are being developed (Ziebart and Dare, 2001 and Ziebart et al, 2002) can cope with highly complex spacecraft structures, and there is no requirement to simplify either the spacecraft structural data, or the derived models. They are applicable at the stages of mission design, operation or scientific analysis, and are designed to be applicable principally to low earth orbiters and medium earth orbit navigation satellites. Models for GLONASS and JASON-1 have been published and the techniques are currently being applied to GPS Block IIR, and a number of other earth observation missions.

- **Proudman Oceanographic Laboratory, <http://www.pol.ac.uk>**

**Satellite orbit determination:** POL has continued to operate a Precise Range and Range-Rate (PRARE) ground-station at Ascension Island as part of the global tracking network for ERS-2.

**Space gravity:** POL has been closely involved in the Science Working Teams for the GRACE and GOCE gravity missions. At the time of writing, GRACE is providing the first data sets of temporal changes in the Earth's gravity field, and a UK consortium including POL, Newcastle and Reading Universities has been assembled to analyse the data. POL has produced a global barotropic ocean model to help remove the effect of high frequency mass movement which will alias the GRACE retrievals. The model has been tested against deep ocean pressure measurements, and compared favourably with a higher resolution, baroclinic model for these high frequency effects. A massively parallel version of the model which incorporates Greens functions to take account of ocean loading and self attraction is currently being tested.

In 1999 GOCE was accepted by the European space Agency for launch in 2006, and planning has since continued apace. GOCE will provide a definitive snapshot of the gravity field and geoid for oceanographic and geodetic studies. Attention has also begun to be addressed to the scientific needs of the generation of space gravity satellites which will follow GRACE and GOCE.

## 8 GRAVITY SURVEYS

- **Proudman Oceanographic Laboratory**, <http://www.pol.ac.uk>

**Absolute gravity:** POL is using the FG5 absolute gravimeter FG5-103 to make measurements of vertical crustal movements near UK tide gauges with long mean sea level records. This is part of a UK program to use GPS and absolute gravity measurements to separate climate related changes in mean sea levels from the vertical crustal component in relative sea levels. Absolute gravity near 3 of the United Kingdom's (UK) core tide gauges was repeatedly measured using an FG5 gravimeter over a period of 3-4 years to determine vertical land movements at those gauges. The absolute gravity sites were established at Newlyn and Aberdeen in 1995 and in Lerwick in 1996. Assuming a height change of 1 mm causes a change in gravity of 0.2 microgal we see vertical land movements of  $1.0 \pm 1.4$  mm/yr at Newlyn,  $-3.8 \pm 1.6$  mm/yr at Lerwick and  $0.9 \pm 3.1$  mm/yr at Aberdeen. These land movements are, within the error estimates, in agreement with land movement predicted by a model of post-glacial rebound/subsidence. To ensure that our absolute gravimeter is giving accurate results with an accuracy equivalent to the highest international standards it has been regularly inter-compared with other absolute gravimeters at sites in USA (1999) and France (2001, ICAG2001) and was verified to be in agreement at the 2 microgal level.

- **United Kingdom Hydrographic Office (UKHO)**, <http://www.ukho.gov.uk>

**OSGM02 model:** The UK Hydrographic Office has provided offshore gravity data in support of the OSGM02 geoid model.

## 9 THEORETICAL GEODESY, EARTH TIDES, EARTH ROTATION AND MISCELLANEOUS GRAVIMETRIC STUDIES

- **School of Civil Engineering and Geosciences, The University of Newcastle**, <http://www.ceg.ncl.ac.uk/>

**Ocean tide loading:** We are currently conducting research into the effects of ocean tide loading in the UK. GPS data collected at permanent sites around the UK are being analysed with the GIPSY/OASIS II software, estimating harmonic terms along with the site coordinates and other parameters. Ongoing research will compare these measurements with estimates from numerical tide models and absolute gravity. A similar project is also underway for the Antarctic continent where large uncertainties exist in numerical tide models, particularly under and near to the ice shelves.

**IGS GNAAC:** The IGS supports the use of GPS by the provision of high quality orbits, Earth rotation parameters and coordinate time series. Newcastle University is a global network associate analysis centre (GNAAC) of the IGS. As such we play a key role in this service by integrating weekly coordinate and Earth Orientation Parameter (EOP) solutions from IGS analysis centres into a common reference frame, for all GPS tracking sites that are a part of the IGS. The TANYA software used for this has been developed at Newcastle, as well as software for online checking of data file compliance to the SINEX format. Global station coordinate and EOP estimates from satellite laser ranging measurements are also being combined as part of the ILRS Pilot Project. The results from these combinations are also of great importance for global plate kinematics studies (*Lavallée, 2000*).

The coordinate time series from the GNAAC has allowed the experimental discovery of a degree-1 global deformation due to seasonal hydrological movements (*Blewitt et al., 2001*) and further developments in the areas of self-consistent loading theory, ocean/continent water distribution, and sea-level change (*Blewitt & Clarke, to be published in 2003*). These data are also currently

being used to validate surface deformations inferred from measurements and models of the time-varying gravity field.

- **Grant Institute, The University of Edinburgh,**  
<http://www.glg.ed.ac.uk/gi/>

#### **Theory (Geoid and Vertical Reference Systems):**

1. *Hipkin, (to be published in 2003)* describes a new and much simpler way to compute the geoid from gravity data available on the surface of an *ellipsoidal* Earth. The conventional ellipsoidal modification of Stokes integral over a local cap misses up to 98% of the ellipsoidal effect - contributions up to 0.6 m - but is also an unnecessary procedure. We only need to compute a correctly ellipsoidal version of the free air anomaly from a global gravity model like EGM96 when using it in a remove-restore algorithm.
2. *Hipkin (2002a)* argues that the geoid can no longer be defined adequately by its relation to mean sea level but that a conventional global datum should be set by giving the total geopotential on the geoid  $W_0$  the same value as the reference potential on the ellipsoid,  $U_0$ . If this conventional datum is adopted and the 1983 IAG resolution about zero-frequency tides is implemented by all, no parameter changes to the Geodetic Reference System 1980 are needed.
3. The theory and practice of a modern vertical reference system with a global datum and no monuments is discussed in *Hipkin (to be published in 2003)*. The essence of the European Vertical Reference System 2000 adheres to this model but its frame is inconsistent by retaining NAP as a definitive datum rather than a provisional 'working hypothesis'.

#### **Theory (Geodynamics):**

1. An analysis (*Hipkin, 2001a,b*) of the statistical properties of data distributed randomly over a sphere concluded that the definitions of the 'power spectrum' adopted in the geodetic and geomagnetic communities are both wrong. Because of the error, a first order property of the Earth's gravity field had been overlooked. With the new analysis, it is shown that the complexity of Earth structure must increase by one or two orders of magnitude between the uppermost mantle and the core, in contrast to the 'empty mantle' models derived from some seismic tomography studies. It is also shown that Moritz' empirical covariance function  $C_3$  is a necessary result of randomly distributed density anomalies over the Bjerhammer sphere but that his alternative  $C_2$  has no such basis in statistical physics.
2. A further application of the pink noise model (*Hipkin, 2002b*) shows its superiority over Kaula's rule as a statistical model of the Earth's gravity field. The systematic deviations of EGM96 from the power law relation of Kaula's rule argue against a 'criticality' as a mechanism for internal density heterogeneities. A nine parameter pink noise model predicts all 120000 free parameters of EGM96 with residuals that are indistinguishable from the normal distribution.
3. An extension of the pink noise model to the gravity fields of the other terrestrial planets (*Hipkin, to be published in 2003*) revealed a quite unexpected identity in the rheological properties of the lithospheres of the Earth and Venus, despite their very different thermal and tectonic regimes. The analysis for the Earth gave a central depth of 49 km and a load-bearing capacity  $10^{4.65}$  Pa, compared with 51 km and  $10^{4.56}$  Pa for Venus. In addition, the larger load-bearing capacity of Venus at 500 km depth is not reached inside the Earth until depths two or three times greater, consistent with the hypothesis the re-circulation of volatiles into the Earth's upper mantle by plate tectonics weakens it substantially compared with Venus, where subduction does not occur.
4. Forsyth's model for interpreting isostatic admittance data in terms of a combination of bottom and top loading has been generalised (Banks et al, 2001) to include internal loading. We

describe an objective way to combine admittance and coherence information that puts unique bounds on rheological properties.

- **Proudman Oceanographic Laboratory**, <http://www.pol.ac.uk>

**Earth tides and ocean tide loading:** Major progress has been made with the interpretation of tidal gravity measurements taken by various groups around the world using continuously recording superconducting gravimeters at a network of global sites as part of the 6 year (1997-2003) Global Geodynamics Project (GGP). In addition, measurements made by POL in the 1980s in Europe, China and Brazil using LaCoste and Romberg ET spring gravimeters with electrostatic feedback proved to be of great value in this work. Ocean tide loading model computations were made for the gravimeter sites using 10 new ocean tide models. It was shown that a few of the superconducting gravimeters have significant calibration errors and are therefore not suitable for testing the latest body tide and ocean tide models. The relatively large number of high quality observations from both superconducting and spring gravimeters in Europe, together with the very small ocean tide loading at diurnal periods in this area, provide a valuable opportunity for testing the models of the Earth's body tide. It was shown that the observed amplitudes are consistent with the latest body tide models and that there is a phase lag of order  $0.02^\circ$  in the diurnal gravity body tide (after correction for ocean tide loading). This is first time that this very small phase lag (due to the Earth's inelasticity) has been observed in tidal gravity measurements.

The global tidal gravity observations were used to test the 10 different ocean tide models. Some of the ocean tide models give anomalous results in various parts of the world. For example, the 4 French global finite element ocean tide models give anomalous results in the western Pacific (China, Japan and Australia). Tidal gravity observations from Spitzbergen provided a valuable opportunity to test the global ocean tide models in the Arctic Ocean, where the models cannot be constrained by Topex/Poseidon altimetry data. It was shown that in this area the latest of the French global finite element ocean tide models is in very close agreement with the tidal gravity measurements and that the agreement is better than that obtained using regional ocean tide models.

## 10 GEOID DETERMINATION

- **Department of Geomatic Engineering, University College London**  
<http://www.ge.ucl.ac.uk/>

**Geoid determination:** Work has been carried out at by the Department of Geomatic Engineering at UCL on geoid determination at local and regional scales. At the local scale, a geoid model was constructed for engineering purposes and compared with OSGM91 (*Iliffe et al, 2000*). More recently, UCL formed a consortium with KMS and the University of Copenhagen in Denmark to compute OSGM02, a new geoid model for the British Isles (*Forsberg et al, 2002*). Full results are to be published shortly.

- **Grant Institute, The University of Edinburgh**,  
<http://www.glg.ed.ac.uk/gi>

### **Practical geoid computation and interpretation:**

1. A new geoid EDIN2000 (*Hipkin et al, to be published in 2003*) has been computed covering the British Isles and adjacent parts of the north west European shelf, specifically for the purpose of determining dynamic sea surface topography over the oceanic parts of the area (*Haines et al, 2003*). Comparisons are made with GPS and levelling, and with oceanographic and gravimetric determinations of dynamic sea surface topography. After decades of controversy, we finally confirm the suspected large systematic error in British levelling, but also find a small local defect in all current geoid models (EGG97, EDIN2000 and OSGM02)

near the Newlyn datum. Probably for the first time globally, we have been able to use the combination of a gravimetric geoid and satellite altimetry to determine realistic geostrophic currents in an oceanic region. We find a three-year mean speed of  $0.32 \text{ m s}^{-1}$  for the shelf edge current carrying 'Gulf Stream' water into the Norwegian Sea, with a noise level elsewhere generally below  $0.05 \text{ m s}^{-1}$ .

2. There is a very significant improvement in the quality of the geoid computed from marine gravity data when the datum and trend of along-track gravity profiles are adjusted to minimise cross-over errors. But we also found that simple interpolation of point values to estimate gravity at the point where two tracks crossed aliases random noise into long-wavelength geoid errors. Consequently a new algorithm has been developed for network adjustment of marine gravity data. It is capable of handling very large numbers of irregularly distributed ship tracks. A pilot study covering all data in the North Sea was presented at IGGeC (*Hipkin, 2002c*) but the method is now being applied to a compilation of 13000 ship tracks covering the whole of the northern North Atlantic as part of the GOCINA project (*Knudsen et al, to be published in 2003*).

- **IESSG, The University of Nottingham**, <http://www.nottingham.ac.uk/iessg/>

**GPS heighting of fundamental benchmarks for geoid determination:** In early 1999, the Ordnance Survey of Great Britain (OSGB) carried out GPS observations at a network of 188 fundamental benchmarks (FBMs), in order to compile a high quality GPS/levelling data set with which to evaluate geoid models for the UK. The data was observed as a series of relatively short, 4 hour observation sessions, with each FBM station being observed for two sessions. The IESSG were responsible for the processing and analysis of the GPS data in order to compute ellipsoidal heights in the ETRS89 e1989.0 coordinate reference system. The results showed that by using 4 hours of data collected with a single GPS receiver and data from active stations that form part of the National GPS Network of Great Britain, it was possible to determine ellipsoidal heights to a precision and accuracy of about 1.5cm. The ellipsoidal heights were used in the computation of the OSGM02.

- **Ordnance Survey of Great Britain**, <http://www.ordsvy.gov.uk>  
**Ordnance Survey Ireland**, <http://www.osi.ie/>  
**Ordnance Survey of Northern Ireland**, <http://www.osni.gov.uk/>

**Geoid Model OSGM02:** Commissioned by the Ordnance Surveys of Great Britain, Ireland and Northern Ireland, the Ordnance Survey Geoid Model 2002 (OSGM02) utilizes all available gravity, terrain and GPS / levelling data and covers the area  $45.5^\circ \text{ N} - 61.5^\circ$  and  $11.5^\circ \text{ E} - 3.5^\circ \text{ W}$  (approximately  $1445 \text{ km} \times 980 \text{ km}$ ). A quasi-geoid was constructed and, subsequently, converted to geoidal heights. This gravimetric geoid was then fitted to GPS / levelling data. To accommodate different vertical datums, the GPS fitting has been computed in patches corresponding to the various datums in use (i.e. Newlyn, Belfast, Malin Head and various island datums) and is limited to within 10 km offshore. The final post-fit standard error of the fitted geoid is 2 cm for mainland Great Britain and Northern Ireland, slightly poorer in the other areas.

A scientific version of the gravimetric geoid, not fitted to the levelling datums, will be made available upon request.

## 11 DEFORMATION MONITORING

- **School of Civil Engineering and Geosciences, The University of Newcastle,**  
<http://www.ceg.ncl.ac.uk/>

**Tectonic deformation monitoring:** We have continued our geodetic investigation of present-day tectonic deformation throughout Greece and the Aegean Sea, assimilating new GPS campaign measurements with previous data (*SING Working Group, 2001*). More recently, we have taken part in large-scale GPS campaigns in mainland Italy and their comparison with historical terrestrial surveys (*Hunstad et al., to be published in 2003*).

Permanent surface displacement due to an individual earthquake is non-linearly related to the earthquake source parameters. This has implications both for the estimation of source parameters from the inversion of geodetic data, and for realistic assessment of the error budget. We have applied robust non-linear inversion methods to recent earthquakes (e.g. *Wright et al., 1999*) to recover realistic source parameters with minimal use of *a priori* information. Recent and continuing work shows that the spatial correlation of geodetic errors is highly significant for error estimation, particularly when using synthetic aperture radar (SAR) interferometry (InSAR).

We have carried out tests of rapid-static, kinematic and real-time-kinematic GPS methods for use in volcano deformation monitoring on Mt Etna, Sicily (*Page, 2000*). Ongoing work includes the use of small-scale GPS networks to mitigate tropospheric errors in InSAR and kinematic GPS, for use in volcano monitoring projects and investigations into the effects of tropospheric estimation in mountainous areas and are developing models to allow improved estimation for RTK heighting in mountainous terrain.

**Local/industrial deformation and subsidence monitoring:** Since 1996 we have been investigating the use of high precision GPS geodesy to monitor platform subsidence rates at several North Sea oil platforms operated by Shell Petroleum and Exploration (UK). The precise point positioning (PPP) strategy is employed as implemented in the GIPSY/OASIS II software developed by JPL. Further research includes investigation of the source of non-secular signals present in these and related time series.

Over the past two years ground subsidence due to oil and mineral extraction has been measured at a number of locations. In the North-East of England, ongoing subsidence due to past coal mining has been monitored using a newly developed low-cost 'all-in-a-box' single-frequency GPS system (*Gledan and Edwards, 2001*). Coastal erosion in North Yorkshire has also been monitored using GPS and InSAR in conjunction with photogrammetric techniques (*Buckley, to be published in 2003*).

- **IESSG, The University of Nottingham,** <http://www.nottingham.ac.uk/iessg/>

**Monitoring changes in regional ground level in the Thames Estuary and Greater London:** Small ground movements can have major strategic or economic significance in certain regions of the world. For example, in low lying river estuaries and coastal regions susceptible to flooding. One such region where small ground movements are exhibited is the Thames Estuary and Greater London, warranting the construction of tidal defences, such as the Thames Barrier. In 1996, the UK Environment Agency initiated a project involving the IESSG and the British Geological Survey. The objective was to develop a generic strategy for monitoring changes in regional ground level using GPS and the ITRS, and providing an interpretation of such changes in terms of local and regional geology. The strategy developed was based around the use of a small number of continuous GPS stations, acting as reference stations for a dense network of monitoring stations, observed using episodic GPS measurements every three months. In a pilot study completed in 1999, 2.25 years of CGPS data and nine sets of episodic GPS measurements were measured at a network of three CGPS stations and twenty-two monitoring stations in the Thames region. The pilot study rationalised the understanding of current changes in ground level in the Thames region. Specifically, over the 2.25 year monitoring period, it was shown that seasonal changes of up to 50 mm occurred where London Clay is at the surface, but such movements were not apparent where London Clay is overlaid by other deposits.



## 12 MEAN SEA LEVEL STUDIES

- **School of Civil Engineering and Geosciences, The University of Newcastle,**  
<http://www.ceg.ncl.ac.uk/>

**Mean sea level studies:** GPS observations at NSTG (see above), along with repeated precise levelling, have been used to calibrate long-term tide gauge measurements of sea-level change. NSTG has a long (100-year) history of tidal observations and is close to a nodal line of post-glacial rebound, making it a key site in climate change studies. However, our GPS and levelling results confirm that the tide gauge structure has significant vertical movement with respect to the nearby bedrock which has no significant geocentric motion (Sanli, 1999; Sanli & Blewitt, 2001).

**IGS/EPN and TIGA stations (MORP and NSTG):** Since 1996, the University of Newcastle has operated a permanent GPS site near Morpeth (MORP), about 30 km north of Newcastle upon Tyne, and carried out episodic and semi-permanent GPS observations at the 100-year-old tide gauge at North Shields (NSTG). In association with the University of Nottingham, NSTG has operated continuously since 2001. In late 2002, MORP was accepted into the IGS and European Permanent Network (EPN), contributing daily and near-real-time data. The entire back-catalogue of GPS data for MORP and NSTG has been submitted to the TIGA archive in support of sea-level change measurements.

- **IESSG, The University of Nottingham,** <http://www.nottingham.ac.uk/iessg/>

**European Sea Level Observing System (EOSS) and European Sea Level Service (ESEAS):** In 1996, European Commission Cost Action 40: EOSS (European Sea Level Observing System) was started. Dr Bingley from the IESSG was UK national delegate to EOSS and was an active member of working group 1 of EOSS, which was specifically concerned with precise height determination at tide gauges. One result of Cost Action 40 was the proposal to establish a European Sea Level Service (ESEAS). Work on the development of ESEAS was started in July 2001 to bring together a major fraction of the previously scattered sea level observing and research resources in Europe into a coordinated research organisation, in order to study sea level variations at inter-annual and century time scales and assess potential future changes in sea level and extreme sea levels. This work was greatly assisted by the award of a three-year European Commission grant through Framework V. The so-called ESEAS-Research Infrastructure (ESEAS-RI) project started in November 2002. The ESEAS-RI project involves twenty-one partners from thirteen countries. It comprises five work packages:

- WP1: Quality control of sea level observations.
- WP2: Absolute sea level variations.
- WP3: Decadal to inter-decadal sea level variations.
- WP4: Improving the sea level observing system.
- WP5: Project management.

The Institute is contributing to work packages WP1, WP4 and WP5, and is leading work package WP2. Work package WP2 will develop the appropriate processing and analysis strategy for the use of continuous GPS (CGPS) at sites close to tide gauges in order to obtain reliable estimates of vertical land movements, and to assess their contribution to changes in relative sea level. The map shows the proposed network of ESEAS Observing Sites, with red dots showing tide gauges that are not co-located with GPS and blue dots showing tide gauges that are co-located with CGPS. Observations from all CGPS stations co-located with ESEAS Observing Sites will be processed and analysed. These sites cover several areas of Europe with types of coasts being affected by different physical processes (e.g. tectonic activity, post-glacial rebound, sedimentary compaction, anthropogenic effects). For further details on ESEAS and EOSS see <http://www.e seas.org/>

**Monitoring of vertical land movements at tide gauge sites in the UK:** Sea level observations at tide gauge sites are corrupted by local, regional or continental vertical land movements. In order to derive estimates of absolute sea level changes, relative tide gauge mean sea level

observations must be corrected for these deformations. This can be carried out by referencing tide gauge benchmarks (TGBMs) to a global reference frame using GPS. The tide gauge (TG) measurements can then be de-coupled from changes in ground level and estimates for the change in absolute sea level obtained. The application of GPS to monitor vertical land movements at selected sites of the UK National Tide Gauge network has been on going at the IESSG since 1990. The work has been funded by the Department for Environment, Fisheries and Rural Affairs (DEFRA) through their long term commission with the Proudman Oceanographic Laboratory (POL). The developments at the IESSG have closely followed the recommendations of the "Carter reports" in 1989 and 1994 and the IGS/PSMSL workshop in 1997. Since the report "United Kingdom Research on Geodesy 1995-1998", several more episodic GPS stations have been upgraded to operate as continuous GPS stations (CGPS) giving a total of seven CGPS@TG stations in the UK. Further stations are currently being planned for the near future to be located on some remote Scottish Islands. A total of thirteen episodic GPS campaigns have now been carried out since 1991, with those in 2002 and 2003 incorporating a quasi-continuous observation strategy for TG sites offering secure operation on a temporary basis. Data from several CGPS@TG stations are now being submitted to the IGS Tide Gauge Benchmark Monitoring Pilot Project (TIGA PP) and to the European Sea Level Service (ESEAS), for which the IESSG operates as one GPS analysis centre.

***Development and evaluation of the dual-CGPS station concept for monitoring vertical station motions at tide gauge sites:*** Vertical station velocities and their associated uncertainties derived from continuous GPS (CGPS) height time series can be used to de-couple secular mean sea level (MSL) trends from vertical land movements at tide gauge sites. In order to precisely describe the often local character of these vertical displacements, the observing CGPS antenna should preferably be as close as possible to the tide gauge itself. This sometimes proves to be difficult, as tide gauges are conveniently situated for tidal observations in port or harbour environments, which can be far from optimal for permanent CGPS installations and observations. In this dual-CGPS station concept, one CGPS station is established at the tide gauge, in order to monitor the local vertical land movements, and a second CGPS station is established on stable rock, further inland within a few kilometres of the tide gauge. The tide gauge CGPS station enables the relative mean sea level to be corrected and an estimate for absolute mean sea level to be obtained. The second CGPS station complements this and helps to describe the underlying geophysical vertical land movements. The dual-CGPS station concept is based on the approach that systematic effects common to both stations, e.g. common periodic signals, can be removed by differencing. The coordinate difference time series derived from the dual-CGPS station analysis show a similar day-to-day scatter as the single baseline time series obtained from the single baseline analysis between the station pair. The findings from the evaluation of the dual-CGPS station concept support the investigations into vertical land movements. Furthermore, the dual-CGPS station concept enables a range of additional methods to be applied for analysis, such as adaptive filtering.

***Stochastic characterisation of the CGPS coordinate time series in the UK:*** The coordinate time series of 21 CGPS stations in the UK have been analysed for their stochastic properties. The coordinate time series have an observation time span of between 1 and 5 years and the data have been archived in the British Isles GPS archive Facility. The assumption that CGPS measurements would improve estimates of station velocities by a factor of  $(1/\sqrt{N})$ , with  $N$  being the number of measurements, by having more measurements than when observing episodically, has been shown to be unrealistic. Station velocity uncertainties in CGPS coordinate time series have been shown to be too optimistic, if the day-to-day scatter in daily coordinate solutions is wrongly assumed to be time independent (white noise). The stochastic characterisation of the CGPS coordinate time series was carried out using two empirical methods and the maximum likelihood estimation (MLE) with integer and fractional spectral indices. The results from these methods agree well and confirmed that the noise in the UK CGPS coordinate time series obtained from the analysis at the IESSG, is not purely white. It was demonstrated that the best stochastic model describing the coordinate time series consists of a combination of white and coloured noise. Using estimates for each noise component it is possible to compute more realistic station velocity uncertainties.

- **Proudman Oceanographic Laboratory**, <http://www.pol.ac.uk>

**Sea level studies:** MSL data from the global network have continued to be collected at POL by the Permanent Service for Mean Sea Level (PSMSL) of the International Council for Science (ICSU). The PSMSL is one of the main organisers of the Global Sea Level Observing System (GLOSS) of the Intergovernmental Oceanographic Commission (IOC). MSL studies have included research into long term changes of UK, European and global MSL, of which the IPCC Third Assessment Report in 2001 was a special focus. Sea level change investigations have also taken place within programmes of the EU and DEFRA, and as part of Programme 1 of the POL Science Plan 2001-6.

Research highlights have included determination of the regional trends in MSL through the 20th century, together with the magnitude of MSL 'accelerations' from the 18th to 20th centuries obtained from 'data archaeology' of some of the first UK sea level information. The role of the North Atlantic Oscillation on UK MSL and wave climate has been investigated at POL through Tyndall Centre collaborations.

MSL information from tide gauges have been demonstrated to be essential to the monitoring of ocean circulation in many ocean areas (e.g. through straits and at high latitudes), often in combination with altimeter information.

POL has continued the long (now 15 year) time series of ocean bottom pressure measurements in Drake Passage, along with tide gauge measurements at Rothera and (in conjunction with the Ukraine) at Vernadsky. An analysis, in conjunction with other tide gauge records from around Antarctica obtained by various international groups as a contribution to GLOSS, has confirmed the model prediction that sea level fluctuations are highly coherent around Antarctica, and are correlated with the atmospheric Southern Hemisphere Annular Mode. The spatial pattern of this coherent mode has been mapped in conjunction with satellite altimeter data. Other relationships between tide gauge data, bottom pressure, bottom temperature, and El Nino, and the sudden change in J2 described in 2002 from satellite orbital computations, are under investigation.

### 13 GEOPHYSICAL AND OCEANOGRAPHIC APPLICATIONS OF GNSS

- **British Antarctic Survey**; <http://www.nerc-bas.ac.uk>

The British Antarctic Survey has been involved in the compilation of three major mapping products:

1. BEDMAP - A gridded compilation of bedrock topography of Antarctica, compiled using a wide variety of international sources (*Lythe et al., 2000*). For more information see <http://www.nerc-bas.ac.uk/public/aedc/bedmap/bedmap.html>
2. ADMAP - A gridded compilation of magnetic anomaly data for Antarctica, compiled using a wide variety of international sources (*Golynsky et al., 2001*). For more information see [http://www.antarctica.ac.uk/mdms/cgi-bin/view.pl?pd\\_pkey=94224675324251](http://www.antarctica.ac.uk/mdms/cgi-bin/view.pl?pd_pkey=94224675324251)
3. The Antarctic Digital Database version 4 - Vector topographic data for the entire continent of Antarctica, managed on behalf of the Scientific Committee on Antarctic Research by British Antarctic Survey. Downloadable from [http://www.nerc-bas.ac.uk/magic/add\\_home.html](http://www.nerc-bas.ac.uk/magic/add_home.html) (*ADD Consortium, 2000*). Latest version is version 4.0, released in August 2002. Version 4.1 due for release in summer 2003.

- **School of Civil Engineering and Geosciences, The University of Newcastle,**  
<http://www.ceg.ncl.ac.uk/>

**Glaciological application of GPS:** We have pioneered a technique that produces highly precise kinematic estimates of horizontal and vertical ice motion using Precise Point Positioning (*King & Aoki, to be published in 2003*). Horizontal coordinate estimates are precise at the 20-30mm level, while vertical coordinates are precise at the 50mm level. Such high precision is obtainable since ice motion is normally slow and constant and hence tight constraints can be applied to the site random walk. This technique has been utilised in research into non-linear horizontal ice stream and ice shelf motion in Antarctica and vertical tidal measurements on several ice shelves. Further work has been done on the Inverse Barometer Effect over the ice shelves. These results have had a significant impact on the modelling of the Antarctic ocean tides and led to the discovery of stick-slip motion in the mouth of Whillans Ice Stream (*Doake et al., 2002*).

- **United Kingdom Hydrographic Office (UKHO),** <http://www.ukho.gov.uk>

**Integrated Coastal Zone Mapping ICZM® Project:** The UKHO has been actively involved with the Ordnance Survey and British Geological Survey on this project. Particular focus is now on the assimilation of vertical datums and further research continues on this subject.

## 14 LAW OF THE SEA

- **United Kingdom Hydrographic Office (UKHO),** <http://www.ukho.gov.uk>

**Law of the Sea:** The UKHO is actively involved with ABLOS, the joint IHO/IAG/IOC Advisory Board on the Law of the Sea. The UKHO provided the Chairman from 1999 – 2001.

## BIBLIOGRAPHY

**Adams J C; Smith M J; Bingley R M;** Terrestrial laser scanning and close-range photogrammetry for cliff-face mapping, *Remote Sensing and Photogrammetry Society 1st Annual Meeting, London, 12-14 September 2001.*

**ADD Consortium;** Antarctic Digital Database, Version 3.0. Database, manual and bibliography, *Scientific Committee on Antarctic Research, Cambridge. Manual and bibliography, 93 pp, 2000.*

**Appleby G M; Otsubo T;** Evaluation of potential systematic bias in GNSS orbital solutions, *Proceedings of 13<sup>th</sup> International Workshop on Laser Ranging, Washington DC, October 2002.*

**Appleby G M; Wood R;** Report on European Workshop at Herstmonceux on Detecting and eliminating errors in EUROLAS network, *ILRS, March 2002.*

**Appleby G M; Otsubo T;** Comparisons of SLR Observations and Orbits with GLONASS and GPS microwave Orbits, *Proceedings of 12<sup>th</sup> Int. Workshop on Laser Ranging, Matera, Italy, November 2000.*

**Appleby G M; Sherwood R A; Rooney E;** The accuracy of GLONASS Broadcast Orbits, *MIT Conference Proceedings, October 2000.*

**Appleby G M; Otsubo T; Sinclair A T;** Comparison of precise SLR orbits of the GLONASS satellites with microwave orbits, *Proceedings of International GLONASS Experiment Workshop, Nashville, Tennessee, Jet Propulsion Laboratory, Pasadena, California, 1999.*

**Appleby G M; Gibbs P; Sherwood R A; Wood R;** Achieving and maintaining sub-cm accuracy for the Herstmonceux single-photon SLR Facility, *Laser Radar Ranging and Atmospheric Lidar Techniques II, SPIE Vol. 3865, 52-63, 1999.*

**Aquino M; Waugh S; Moore T; Dodson A H;** Testing the Performance of the WBMOD Scintillation Prediction Model against GPS Scintillation Data from the November 2001 Storm, *IES 2002 - Ionospheric Effects Symposium 2002, Alexandria, VA – USA, 7 - 9 May 2002.*

**Aquino M; Moore T; Dodson A H; Waugh S;** Ionospheric Scintillation Effects on GPS Measurements, *invited presentation to the Radiocommunications Agency's Ionospheric Effects on Radio Systems Task Group Meeting, London, 21 November 2002.*

**Aquino M H O; Waugh S; Dodson A H; Moore T; Skone S;** GPS based ionospheric scintillation monitoring, *Proceedings of Space Weather Workshop: Looking towards a European Space Weather Programme, 6 pages, ESTEC, Noordwijk, The Netherlands, 17-19 December 2001.*

**Aquino M H O; Dodson A H; Bingley R M; Penna N T;** The Establishment of a UK National Network of Continuously Operating GPS Receivers (COGRs), *Proc International Symposium on GPS : GPS99, Tsukuba, Japan, October 1999.*

**Aquino M H O; Ashkenazi V; Moore T;** Prospective Use of Wide Area Augmentations in Brazil, *Proc GIS Brasil 99, V Latin American Congress for Users of Geoprocessing, Salvador, Brazil, July 1999.*

**Archer S; Gikas V; Pinel C; Ridyard D; Cross P;** Spatially and temporally correlated navigation errors: how do they manifest themselves in seismic data?, *First Break, Vol 17, No 11, pp355-362, 1999.*

**Ashkenazi V;** Directions 2002, *GPS World Showcase, Vol 12, No 12, p11, December 2001*

**Ashkenazi V;** Satellite Navigation and Positioning: State of the Art, *Proc Geomark 2000, ESA SP-458, ISBN 92-9092-673-2, Paris, France, April 2000.*

**Ashkenazi V;** Synergy Between SATNAV, SATCOM and EO: State-of-the-Art 2000-2015, *Proc Geomark 2000, ESA SP-458, ISBN 92-9092-673-2, Paris, France, April 2000.*

**Ashkenazi V;** Galileo: The Challenge, *Proc GNSS 2000, 4th European Conference on Global Navigation Satellite Systems, Edinburgh, May 2000.*

**Ashkenazi V;** The Challenges facing Galileo, *Space Policy, Vol 16, pp185 - 188, August 2000.*

**Ashkenazi V;** Directions 2001, *GPS World Showcase, Vol 11, No 12, ISSN 1048-5104, 2000.*

**Ashkenazi V;** Ubiquitous Computing Infomobility and the Role of Navigation, *Proc Industrial Workshop on Emerging Satellite Navigation Markets, Vienna, Austria, November 2000.*

**Ashkenazi V; Dumville M;** GPS, Galileo and Wireless Technologies, *World Mobile Technology, Cornhill Publications Ltd, Issue 3, pp 84 - 85, 2000.*

**Ashkenazi V; Dumville M; Park D;** Robot Positioning and the Global Navigation Satellite System, *Industrial Robot Journal, Vol 27, No 6, 2000.*

**Ashkenazi V; Bingley R M; Dodson A H; Booth S; Penna N T; Baker T F; Greenaway R; Nursey K; Bedlington D; Arthurton R; Ellison R;** Monitoring vertical land movements by GPS for flood and coastal protection, 3.4.1 - 3.4.12 in, *MAFF Conference of River and Coastal Engineers, Keele University, 1 - 3 July 1998, MAFF, Flood and Coastal Defence with Emergencies Division, 1999.*

**Ashkenazi V;** Contribution to Directions 2000, *GPS World Showcase, Vol 10, No 12, ISSN 1048-5104, pp 46, December 1999.*

**Ashkenazi V;** Galileo: Challenge and Opportunity, *Galileo's World, Vol 1, No 1, pp 42-44, Winter 1999.*

**Ashkenazi V;** State of the Art in the Synergetic Use of Satellite Observations, *Proc Fourth SAI Annual Users' Seminar, Baveno, Italy, May 1999.*

**Ashkenazi V;** Implementation of WGS 84 for Civil Aviation in Preparation for Satellite Navigation in Europe, *Proc Research Seminar in the Università degli Studi di Bologna, Bologna, Italy, June 1999.*

**Ashkenazi V;** Satellite Navigation: Current Systems, *Proc ASTRON Workshop, Ispra, Italy, July 1999.*

**Ashkenazi V;** GNSS-1 and GNSS-2, *Proc ASTRON Workshop, Ispra, Italy, July 1999.*

**Ashkenazi V;** Applications and Markets, *Proc ASTRON Workshop, Ispra, Italy, July 1999.*

**Ashkenazi V;** Galileo : A New European Satellite Navigation System, *Proc GIS Brasil 99, V Latin American Congress for Users of Geoprocessing, Salvador, Brazil, July 1999.*

**Ashkenazi V;** Combining Earth Observation, Satellite Navigation and Satellite Communications, *Proc GIS Brasil 99, V Latin American Congress for Users of Geoprocessing, Salvador, Brazil, July 1999.*

**Ashkenazi V; Berthelot H;** GPS to Galileo: A European Path, *Air and Space Europe, Vol 1, No 3, 1290-0958, pp 66-71, May / June 1999.*

**Ashkenazi V; Dumville M; Moore T; Roberts G W; Zhang K F; Close G; Lee C; Moore R; Tindall I;** Real Time River Level Monitoring System Based on GPS, *Proc GIS Brasil 99, V Latin American Congress for Users of Geoprocessing, Salvador, Brazil, July 1999.*

**Ashkenazi V; Moore T; Dumville M; Chen W; McPherson D; Millen K; Greenland A; Ward, N; Savill, M; Ruttle, S;** GNSS & WGS 84 for Marine Navigation in UK Waters, *Journal of Navigation, Vol 52, No 2, ISSN 0373-4633, pp 189 - 195, May 1999.*

**Ashkenazi V; Moore T; Dumville M; Chen W; McPherson D; Simpson D; Greenland A; Ward N; Savill M; Ruttle S;** A Realisation of WGS 84 and GNSS for Marine Navigation in the UK, *Proc ION-GPS-99, The 12th International Technical Meeting of the Satellite Division of the Institute of Navigation, 8 pages, Nashville, USA, September 1999.*

**Ashkenazi V; Moore T; Dumville M; Chen W; McPherson D; Simpson D; Greenland A; Ward N; Savill M; Ruttle S;** Practical Considerations for the Introduction of GNSS and WGS 84 for Marine Navigation, *Proc GNSS 99, 3rd European Symp on Global Navigation Satellite Systems, pp 519-524, Genoa, Italy, October 1999.*

**Ashkenazi V; Moore T; Dumville M; Chen W; McPherson D; Simpson D; Greenland A; Ward N; Savill M; Ruttle S;** GNSS and WGS 84 for Marine Navigation, *Proc NAV 99 / ILA 28, International Conference of the Royal Institute of Navigation and the International Loran Association, pp 39-1-39-9, London, November 1999.*

**Baker D F; Dodson A H; Moore T;** Olympic Class GNSS: Mapping Skeleton Sled Tracks, *Galileo's World, Vol 4, No 1, ISSN 1528-2341, pp 10 - 20, Spring 2002.*

**Baker D F;** The Performance of Hybrid GPS and GLONASS, *PhD thesis, University of Nottingham, 2001.*

**Baker H C; Dodson A H; Penna N T; Higgins M; Offiler D;** Ground-based GPS water vapour estimation for meteorological forecasting, *J. Atmos & Solar Terrestrial Phys, 63, pp 1305-1314, 2001.*

**Baker H C; Dodson A H; Penna N T; Higgins M; Offiler D;** Ground-based GPS Water Vapour Estimation: Potential for Meteorological Forecasting, *Submitted to the Journal Atmospheric and Terrestrial Physics, July 1999.*

**Baker H C; Dodson A H; Penna N T; Buerki B; Elgered G; Rius A; Rothacher M;** The WAVEFRONT project on ground-based GPS water vapour estimation, *Proc GIS Brasil 99, V Latin American Congress for Users of Geoprocessing, Brazil, July 1999.*

**Baker H C; Dodson A H; Penna N T; Higgins M; Offiler D;** Ground-Based GPS Water Vapour Estimation: Potential for Meteorological Forecasting, *Proc Symposium JSG28, IUGG General Assembly, Birmingham, July 1999.*

**Baker T F; Bingley R M;** GPS monitoring of changes in ground level for flood and coastal defence, *Flood and Coastal Defence Research News (DEFRA/EA), 2: 6-7, 2002.*

**Baker T F; Bingley R M; Teferle F N; Dodson A H; Williams S D P;** GPS and absolute gravity at UK tide gauges, *Proceedings of GLOSS Group of Experts Meeting, IOC, University of Hawaii, Hawaii, April 2001. Intergovernmental Oceanographic Commission Workshop Report, 180, 2002.*

**Baker T F; Bos M S; Williams S D P;** Confronting superconducting and absolute gravity measurements with models, *Bulletin d'inf. Marées Terr.*, 136, 10743-5, 2002.

**Baker T F; Blackman D L; Flather R A; Vassie J M; Woodworth P L;** Integrated effects of climate change on coastal extreme sea levels, *Proceedings of the 36th DEFRA Conference of River and Coastal Engineers, Keele University, 20-22 June 2001, O3.4.1-O3.4.12, 2001.*

**Baker T F; Bos M S;** Tidal gravity observations and ocean tide models, *Journal of the Geodetic Society of Japan*, 47: 76-81, 2001.

**Balmino G; Rummel R; Visser P; Woodworth P L;** Gravity Field and Steady-State Ocean Circulation Mission, Reports for assessment: the four candidate Earth Explorer Core Missions, *European Space Agency Report SP-1233(1). 217pp., 1999.*

**Banks R J; Roger J; Francis S C; Hipkin R G;** Effect of loads in the upper crust on estimates of the elastic thickness of the lithosphere, *Geophysical Journal International*, 145, 291-299, 2001.

**Barnes J B;** Real time kinematic GPS and multipath : characterisation and improved least squares modelling, *Ph.D. thesis, University of Newcastle upon Tyne, 2000.*

**Becker M; Zerbini S; Baker T F; Burki B; Galinis J; Garate J; Georgiev I; Kahle H; Kotzev V; Lobazov V; Marson I; Negusini M; Richter B; Veis G; Yuzefovich P;** Assessment of height variations by GPS at Mediterranean and Black Sea coast tide gauges from the SELF project, *Global and Planetary Change*, 34, 5-35, 2002.

**Bennett J; Hilton, R D; Berry P A M;** A simulator for predicting altimeter waveforms over land. *European Geophysical Assembly, Nice France, April 2002.*

**Bennet J; Berry P A M;** Expert system slope validation. *BGAS meeting, Herstmonceux castle, 2001.*

**Bennett J; Berry P A M;** An altimeter simulator for land studies. *BGAS meeting, Met. Office training college Reading, United Kingdom, 1999.*

**Berry P A M;** Global river and lake height monitoring using satellite altimeter data. *Hydropower & Dams, Issue 6, 2002.*

**Berry P A M;** Waveform Retracking. *International Altimeter Workshop, Wuhan, China, August 2002.*

**Berry P A M; Hilton R D; Johnson C P D;** ACE: first full release of the new 30" GDEM incorporating satellite altimeter derived heights. *European Geophysical Assembly, Nice France, April 2002.*

**Berry P A M; Johnson C P D; Pinnock R A;** Natural Land targets for ERS and Envisat backscatter monitoring. *IAG Scientific Assembly September 2001.*

**Berry P A M; Hilton R D; Johnson C P D;** Global Digital elevation models –current and future. *IAG Scientific Assembly September 2001.*

**Berry P A M; Hilton R D; Johnson C P D; Pinnock R A;** The New ACE GDEM - Implications For Gravity Field Models. *European Geophysical Assembly, Nice, France, April 2001.*

**Berry P A M; Pinnock R A;** Use of natural land targets for monitoring ERS-2 and Envisat altimeter sigma0. *European Geophysical Assembly, Nice, France, April 2001.*

**Berry P A M;** Land applications of ERS satellite altimetry. *BGAS meeting, Herstmonceux castle, 2001.*

**Berry P A M; Dowson M; Pinnock R A;** An independent evaluation of accuracy of regional DEMs using satellite altimeter data. *European Geophysical Assembly, Nice, France, April 2000.*

**Berry P A M; Pinnock R A; Hilton R D; Johnson C P D;** A new Global Digital Elevation Model at 30" resolution: first generation. *European Geophysical Assembly, Nice, France, April 2000.*

**Berry P A M; Johnson C P D; Hilton R D;** Correction of Digital elevation data using altimeter derived heights. *European Geophysical Assembly, Nice, France, April 2000.*

**Berry P A M; Pinnock R A; Johnson C P D; Hilton R D; Dowson M;** ACE: a new Global Digital Elevation Model. *IAG General Assembly, Banff, Canada July, 2000.*

**Berry P A M; Pinnock R A; Hilton R D; Johnson C P D; Dowson M;** Characteristic Error Signatures of Current Global Digital Elevation Models. *Poster presentation. IAG General Assembly, Banff, Canada July, 2000.*

**Berry P A M; Pinnock R A; Hilton R D; Johnson C P D;** ACE: a new GDEM incorporating satellite altimeter derived heights. *ERS-ENVISAT symposium "Looking Down to Earth in the New Millennium", Gothenburg, Sweden, October 2000.*

**Berry P A M; Pinnock R A; Wilson H K;** Land calibration and monitoring of Envisat RA-2 sigma0. *ERS-ENVISAT symposium "Looking Down to Earth in the New Millennium", Gothenburg, Sweden, October 2000.*

**Berry P A M; Hoogerboord J E; Pinnock R A;** Land calibration of radar altimeter sigma0. *European Geophysical Assembly, The Hague, Netherlands, April 1999*

**Berry P A M; Hoogerboord J E; Pinnock R A;** Identification of common error signature in Global Digital Elevation Models. *European Geophysical Assembly, The Hague, Netherlands, April 1999.*

**Berry P A M;** Topography from land altimeter data: possibilities and restrictions. *European Geophysical Assembly, The Hague, Netherlands, April 1999.*

**Berry P A M; Pinnock R A; Hoogerboord J E;** Land calibration of altimeter sigma0. *BGAS meeting, Met. Office training college Reading, United Kingdom, 1999.*

**Berry P A M; Hoogerboord J E; Pinnock R A;** Global Digital Elevation Models – fact or fiction? *BGAS meeting, Met. Office training college Reading, United Kingdom, 1999.*

**Berry P A M; Hoogerboord J E; Pinnock R A;** An independent global assessment of the quality of current Global Digital Elevation Models. *IUGG XXII General Assembly, University of Birmingham, United Kingdom, July 1999.*

**Berry P A M; Hoogerboord J E; Pinnock R A;** Natural land targets for altimeter sigma0 calibration. *IUGG XXII General Assembly, University of Birmingham, United Kingdom, July 1999.*

**Berry P A M; Pinnock R A; Hoogerboord J E;** Environmental signals in land altimeter data: potential for soil moisture measurement. *IUGG XXII General Assembly, University of Birmingham, United Kingdom, July 1999.*

**Berry P A M;** Global Digital Elevation Models – Fact or Fiction? *Astronomy and Geophysics Vol. 40, Issue 3, PP3.10-3.13, June 1999.*

**Bingley R M; Dodson A H; Penna N T; Baker T F;** Using a 'GPS/MSL Geoid to test Geoid Models in the UK, Vertical Reference Systems, *Ed. Drewes, H, International Association of Geodesy Symposia, Vol 124, pp 197-202, ISBN 3-540-43011-3, Springer Verlag, 2002.*

**Bingley R M; Dodson A H; Teferle F N; Baker T F;** GPS monitoring of changes in ground level for flood and coastal defence, *Proceedings of 36th Annual Conference of River and Coastal Engineers, Keele University, pp. 03.5.1 – 03.5.12, 2002.*

**Bingley R M; Dodson A H; Penna N T; Teferle F N; Baker T F;** Monitoring the vertical land movement component of changes in mean sea level using GPS: results from tide gauges in the UK, *Journal of Geospatial Engineering, 3: 9-20, 2001.*

**Bingley R M; Dodson A H; Penna N T; Teferle F N; Booth S J; Baker T F;** Using a Combination of Continuous and Episodic GPS data to Separate Crustal Movements and Sea Level Changes at Tide Gauges in the UK, *Proc WEGENER 2000, The Tenth General Assembly of the Working Group of Eu*



**Brown C J; Karuna R; Ashkenazi V; Roberts G W; Evans R;** Monitoring of Structures using GPS, *Proc Institution of Civil Engineers, Structures, ISSN 0965 092X, pp 97 - 105, February 1999.*

**Carleton C;** Co-Author "Developments in the Technical Determination of Maritime Space: Delimitation, Dispute Resolution, Geographical Information Systems and the Role of the Technical Expert", *Maritime Briefing Volume 3 No. 4 published by the International Boundaries Research Unit of Durham University 2002.*

**Carleton C;** Contributing Author "The Razor's Edge" - The Development of Maritime Zones and Boundaries from 1964 to the Present, *published by Kluwer Law International 2002.*

**Carleton C;** Co-Author "Developments in the Technical Determination of Maritime Space: Charts, Datums, Baselines, Maritime Zones and Limits", *Maritime Briefing Volume 3 No. 3, published by the International Boundaries Research Unit of Durham University, 2001.*

**Carleton C;** Member of the Group of Experts, convened by the UN Office for Ocean Affairs and the Law of the Sea in 1999, on Maritime Delimitation, *UN Publication, 2000.*

**Carleton C;** Co-Chief Editor of IOC/IHO book "Continental Shelf Limits: The Scientific and Legal Interface", *published by Oxford University Press, New York, 2000.*

**Carroll J D M;** VideoRoute: an integrated navigation system for rapid mapping, *M.Phil. thesis, University of Newcastle upon Tyne, 1999.*

**Chen W; Celik C T; Bingley R M; Ashkenazi V;** Velocity Field Estimation for Geodetic Networks Using Modified Kalman Filters, *Proc XXII General Assembly of the International Union of Geodesy and Geophysics, Birmingham, July 1999.*

**Church J A.; Gregory J M; Huybrechts P; Kuhn M; Lambeck K; Nhuan M T; Qin D; Woodworth P L;** Changes in sea level. In, *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change. (eds. J.T. Houghton, Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell and C.A. Johnson). Cambridge: Cambridge University Press. 881pp, 2001*

**Close G; Lee C; Moore R; Moore T;** Water Level Monitoring from Space, *Proc Geomark 2000, ESA SP-458, ISBN 92-9092-673-2, Paris, France, April 2000.*

**Clarke P; Cross P A; Cruddace P R; Billiris H; Galanis I; Paradisis D; Veis G; England P; Parsons B; Avalone A; Briole P; Lyon-Caen H; Cocard M; Hollenstein C; Kahle H; Ambrosius B; Stavrakakis G;** Crustal Strain in Greece and the Aegean from Multiply Repeated GPS Measurements in the Interval 1988—2000. *Proceedings of AGU Fall Meeting, San Francisco, America, 2001*

**Clarke P J;** Geodesy and geodetic measurements, in: *The Oxford Companion to the Earth, ed. Hancock PL and Skinner BJ, pp. 398-402, Oxford University Press, 2000.*

**Clarke P J;** Geomatics, *ibid.*, pp.423-424.

**Clarke P J;** Global Positioning System, *ibid.*, pp. 458-459.

**Clarke P J;** Planetary geodesy, *ibid.*, p. 825.

**Clarke P J;** Very long baseline interferometry, *ibid.*, pp. 1087-1088.

**Cross P A; Ziebart M K;** LEO GPS Attitude Determination Algorithm Designed for Real-time On-board Execution, *Proceedings of ION GPS-2002, Portland, Oregon, pp1051-1063, 2002.*

**Cross P A; Keenan R;** Stochastic Modelling: the crucial element in high fidelity DGPS quality assessment?, *Proceedings of HYDRO 2001, Norwich, UK, 2001.*

**Cross P A;** Prospects for GPS - new systems, new applications, new techniques, *Engineering Surveying Showcase, Issue 1, pp10-14, 2000.*

**Cross P A;** Satellite positioning - the future, *Civil Engineering Surveyor December, p14-18, 2000.*

**Cross P A; Higgins M; Lott R;** Reference frames in practice: The role of professional, scientific and commercial organisations, *Proceedings of Quo Vadis Surveying of the 21st Century, FIG Working Week, Prague. 2000*

**Cross P A; Keenan R; Barnes J B; Anthis N; Chagani K; Walker A; Muller J-P; Kitmitto K;** GPS validation of IfSAR digital elevation models from LANDMAP, *Proceedings of the 26th Annual Conference of the Remote Sensing Society, RSS2000, University of Leicester, UK. ISBN 0 946226 296/0 946226 30.X, 2000.*

**Cross P A;** Accuracies achievable by GPS in practical engineering and surveying applications, *Proceedings of Geodesy and Surveying in the Future: The Importance of Heights, National Land Survey of Sweden, Gävle, Sweden, pp. 175-184, 1999.*

**Cruddace P R; Wilson I; Greaves M; Euler H-J; Keenan R; Wuebbena G;** The long road to Establishing a National Network RTK Solution. *Proceedings of the FIG conference Washington DC, 2002.*

**Cruddace P R; Greaves M;** National Report of Great Britain, *Report on the Symposium of the IAG Subcommission for Europe (EUREF) held in Ponta Delgada, 5 – 8 June 2002*, Torres, J. A., Hornik, H. (eds), Bundesamt für Kartographie und Geodäsie, Frankfurt, Germany, 2002. Note – not yet published at time of writing. Expected publication date June 2003. Available on line at [http://www.euref-iaag.org/book2002/5\\_10.PDF](http://www.euref-iaag.org/book2002/5_10.PDF)

**Cruddace P R;** Ordnance Survey's National GPS Network. *Geoinformatics July /August 2001*.

**Cruddace P R;** Britain gets GPS Service via the Web. *Civil Engineering Surveyor, GIS/GPS supplement, 2001*.

**Cruddace P R; Clarke P;** Is GPS at last becoming useful? *BMC Summit magazine, Issue 19, Autumn 2000*

**Cruddace P R; Cross P A; Veis G; Billiris H; Paradissis D; Galanis J; Lyon-Caen H; Briole P, Ambrosius B A C; Simons W J F; Roegies E; Parsons B E, England; Kahle H-J; Cocard M; Yannick P; Stavrakakis G; Clarke P; Lilje M;** An interdisciplinary approach to studying seismic hazard throughout Greece. In *Geodesy Beyond 2000: the Challenges of the First Decade. International Association of Geodesy Symposia, Springer, Vol 121, p279-284, 2000. Also in Destructive Earthquakes: Understanding Crustal Processes Leading to Destructive Earthquakes, Proceedings of The Second EU-Japan Workshop on Seismic Risk. Reykjavik, Iceland, pp98-103, 2000.*

**Cruddace P R; Cross P A;** Overall GPS strain determination and assessment of seismic hazard in Greece. *Dissemination workshop of the EC project SERGISAI and SEISMODOC. Brussels, Belgium, 1999. Springer, 2000.*

**Cruddace P R;** Crustal deformation throughout Greece from a decade of GPS measurements. *American Geophysical Union Fall Meeting. San Francisco, U.S.A. EOS Transactions, v80 no.46 p274, 1999.*

**Cullen R; Moore P;** A spectral optimally interpolated mean sea surface from satellite radar altimetry, *Journal of Geodesy, 75(4), 175-187, 2001.*

**Davies P; Greaves M; Dodson A H; Bingley R M;** National Report of Great Britain, *Report on the Symposium of the IAG Sub-commission for the European Reference Frame (EUREF), Prague, Czech Republic, ISSN 0340-7691. ISBN 3769696220 pp.224225, 1999.*

**Doake C S M; Corr H F J; Nicholls K W; Gaffikin A; Jenkins A; Bertiger W I; King M A;** Tidal movement of Brunt Ice Shelf, Antarctica. *Geophys. Res. Lett., 29(8), doi:10.1029/2001GL014606, 2002.*

**Dodson A H; Chen W; Penna N T; Baker H C;** GPS estimation of atmospheric water vapour from a moving platform, *J. Atmos & Solar-Terrestrial Phys, 63, pp 1331-1341, 2001.*

**Dodson A H; Moore T; Aquino M H O;** Ionospheric Scintillation Effects on GPS Measurements, *Proc GNSS 2001, 5th International Conference on Global Navigation Satellite Systems, Seville, Spain, May 2001.*

**Dodson A H; Meng X; Roberts G W;** Adaptive FIR Filtering for Multipath Mitigation and its Application for Large Structural Deflection Monitoring, *Proc International Symposium on Kinematic Systems in Geodesy, Geomatics and Navigation (KIS 2001), Banff, Canada, 5-8 June 2001.*

**Dodson A H; Moore T; Pattinson M;** Measuring Tropospheric Delay from a Floating Platform, *Proc ION GPS 2001, 14th Technical Meeting of the Satellite Division of the Institute of Navigation, 8 pages, Salt Lake City, September 2001.*

**Dodson A H; Bingley R M; Penna N T; Aquino M H O;** A National Network of Continuously Operating GPS Receivers for the UK, *IAG Symposia Geodesy Beyond 2000, Springer-Verlag, Vol 121, ISBN 3-540-67002-5, pp 367 - 372, 2000.*

**Dodson A H; Moore T;** Geodetic Techniques, *Continental Shelf Limits: The Scientific and Legal Interface, Oxford University Press, ISBN 0-19-511782-4, pp 87 - 104, 2000.*

**Dodson A H; Baker H C; Penna N T; Buerki B; Elgered G; Rius A; Rothacher M;** The WAVEFRONT Project on GPS Water Vapour Estimation, *Pres International Symposium on GPS : GPS 99, Tsukuba, Japan, October 1999.*

**Dodson A H; Baker H C; Penna N T; Higgins M; Jerrett D;** GPS Tropospheric Water Vapour for Meteorological Forecasting, *Proc GNSS 99, 3rd European Symp on Global Navigation Satellite Systems, pp249-254, Genoa, Italy, October 1999.*

**Dodson A H; Bingley R M; Penna N T; Aquino M H O;** A National Network of Continuously Operating GPS Receivers for the UK, *Proc Symposium G1 IUGG General Assembly, Birmingham, July 1999.*

**Dodson A H; Bingley R M; Symons L J; Penna N T;** A National Network of Continuously Operating GPS Receivers (COGRs) for the UK, *Proc XXII General Assembly of the International Union of Geodesy and Geophysics, Birmingham, July 1999.*

**Dodson A H; Chen W; Baker H C; Penna N T; Roberts G W; Jeans R J; Westbrook J;** Assessment of the EGNOS Tropospheric Correction Model, *Proc ION-GPS-99, The 12th International Technical Meeting of the Satellite Division of the Institute of Navigation, 7 pages, Nashville, USA, September 1999.*

**Dodson A H; Chen W; Penna N T; Baker H C;** GPS Estimation of Atmospheric Water Vapour from a Moving Platform, *Proc Symposium JSG28, IUGG General Assembly, Birmingham, July 1999.*

**Dodson A H; Curtis D J; Baker T F;** Ocean Tide Loading Effects on Height, *Proc Geodesy and Surveying in the Future-The Importance of Heights, LMV-Rapport 1999:3, ISSN 0280-5731, pp 261-272, March 1999.*

**Dodson A H; Moon G V; Moore T; Jones D;** Guiding Blind Pedestrians with a Personal Navigation System, *Journal of Navigation, Vol 52, No 3, ISSN 0373-4633, pp 330-341, September 1999.*

**Dodson A H; Moore T;** Geodetic Techniques, Sovereign Limits Beneath the Oceans: Delimiting the New Continental Shelf, *IHO/IOC, In Print 1999.*

**Dodson A H; Moore T; Baker D F; Swann J W;** Hybrid GPS + Glonass, *GPS Solutions, Vol 3(1), ISSN 1080-5370, pp 32-41, July 1999.*

**Dodson A H; Moore T; Moon G V;** A Navigation System for the Blind Pedestrian, *Proc GNSS 99, 3rd European Symp on Global Navigation Satellite Systems, pp 513-518, Genoa, Italy, October 1999.*

**Dodson A H; Penna N T; Baker H C; Higgins M; Offiler D;** The UK GPS IWV Meteorological Forecasting Impact Assessment, *Pres International Symposium on GPS: GPS 99, Tsukuba, Japan, October 1999.*

**Dong X; Woodworth P L; Moore P; Bingley R;** Absolute calibration of the TOPEX/Poseidon altimeters using UK tide gauges, GPS and precise local geoid differences, *Marine Geodesy, 25, 189-294, 2002.*

**Dong X; Woodworth P L; Moore P; Bingley R M; Huang C;** Absolute Calibration of the ERS-2 Altimeter using UK Tide Gauges, *Proceedings of the International Workshop on Satellite Altimetry for Geodesy, Geophysics and Oceanography, Wuhan, China, 8-13 September 2002.*

**Dong X; Woodworth P L; Moore P; Bingley R M;** Absolute Calibration of TOPEX, POSEIDON and JASON-1 using UK Tide Gauges, GPS and a Local Precise Geoid Model, *Proceedings of the Asia Pacific Space Geodynamics (APSG) Sea Level Meeting, University of Hawaii, April 2001.*

**Dowson M; Berry P A M;** A global multi-mission, multi-mode altimeter study in preparation for ENVISAT. *European Geophysical Assembly, Nice France, April 2002.*

**Dowson M; Berry P A M; Bennett J;** Potential impact of ENVISAT on land altimeter coverage. *IAG scientific assembly September 2001.*

**Dowson M; Berry P A M;** Simulator for ENVISAT resolution switching algorithm. *European Geophysical Assembly, Nice France, April 2001.*

**Dowson M; Berry P A M;** ENVISAT RA-2: Potential for land topographic mapping. *ERS-ENVISAT symposium "Looking Down to Earth in the New Millennium", Gothenburg, Sweden, October 2000.*

**Dowson M; Berry P A M;** Verification of altimeter heighting. *BGAS meeting, Met. Office training college Reading, United Kingdom, 1999.*

**Dowson M; Berry P A M; Hoogerboord J E;** An evaluation of techniques for digital elevation model correction using satellite altimeter data. *IUGG XXII General Assembly, University of Birmingham, United Kingdom, July 1999.*

**Dowson M; Berry P A M;** A global precision estimate of radar altimetry heighting over land. *European Geophysical Assembly, The Hague, Netherlands, April 1999.*

**Edwards S J; Cross P A; Barnes J B; Betaille D;** A methodology for benchmarking real time kinematic GPS. *Survey Review. Vol 35, No 273, pp. 163-174, 1999. Reprinted in The Hydrographic Journal, No 94, pp11-15, 1999.*

**Evans A J; Roberts G W; Ashkenazi V;** Improving Position Solutions Over Long Baselines Using IGS, *Proc IAG 2001 Scientific Assembly, Budapest, Hungary, 2-8 September 2001.*

**Evans A J; Roberts G W; Dodson A H; Denby B; Hollands R; Coope, S;** A New Vision for Engineering Geodesy, *Proc IAG 2001 Scientific Assembly, Budapest, Hungary, 2-8 September 2001.*

**Evans A J;** Long Baseline OTF Kinematic GPS, *PhD thesis, University of Nottingham, 2001.*

**Evans A J; Roberts G W; Ashkenazi V;** Precise Kinematic Positioning for Aerial Survey, *Proc GNSS 2000, 4th European Symposium on Global Navigation Satellite Systems, Edinburgh, May 2000.*

**Featherstone W E; Olliver J G;** A Review of Geoid Models over the British Isles: Progress and Proposals, *Survey Review, 36, 280, 78-100, 2001*

**Flather R A; Baker T F; Woodworth P L; Vassie J M; Blackman D L;** Integrated effects of climate change on coastal extreme sea levels, *Proceedings of 36th Annual Conference of River and Coastal Engineers, Keele University, pp 03.4.1 – 03.4.12, 2002.*

**Flather R A; Baker T F; Woodworth P L; Blackman D L;** Integrated effects of climate change on coastal extreme sea levels, *Proudman Oceanographic Laboratory, Internal Document, No 140, 2001.*

**Forsberg R; Strykowski G; Iliffe J C; Ziebart M K; Cross P A; Tscherning C C; Cruddace P; Finch O; Bray C; Stewart K;** OSGM02: A new Geoid Model of the British Isles, *Proceedings of Gravity and Geoid 2002 - GG2002, 3rd meeting of the International Gravity and Geoid Commission, August 26 -30, 2002, Thessaloniki, Greece, 2002. To be published by Springer Verlag in International Association of Geodesy Symposia series, 2002.*

**Gibbs P; Koidl F; Kirchner G;** Range comparison results for various EUROLAS SR timers, *Proceedings of 13<sup>th</sup> International Workshop on Laser Ranging, Washington DC, October 2002*

**Gibbs P; Benham D J; Sherwood R A; Standen P; Walters D; Wood R; Appleby G M;** An overview of Quality Control for the Herstmonceux SLR Station, *Proceedings of 12<sup>th</sup> International Workshop on Laser Ranging, Matera, Italy, November 2000.*

**Gibbs P; Wood R;** C-SPAD as a single-photon detector, *Proceedings of 12<sup>th</sup> International Workshop on Laser Ranging, Matera, Italy, November 2000*

**Gikas V; Cross P A; Ridyard D;** Reliability analysis in dynamic systems; Implications for positioning marine seismic networks, *Geophysics Vol 64, No4, pp1014-1022, 1999.*

**Gille S T; Hughes C W;** Aliasing of high-frequency variability by altimeter observations: Evaluation from bottom pressure recorders, *Geophys. Res. Letters, 28, 1755-1758, 2001.*

**Gledan J; Edwards S J;** GPS for Monitoring Dams and Other Large Structures, *Civil Engineering Surveyor, 22-24, 2001*

**Golynsky A V; Morris P; von Frese R; ADMAP Group;** ADMAP - Magnetic anomaly map of the Antarctic. 1:10,000,000 scale map, *BAS (Misc) 10. Cambridge, British Antarctic Survey, 2001.*

**Greaves M; Cruddace P R;** The OS's new coordinate Transformation for Great Britain. *Geomatics World Volume 10, Issue 1, 2001*

**Greaves M; Cruddace P R;** The Adoption of ETRS89 as the national Mapping System for GB, via a Permanent GPS Network and Definitive Transformation, *Report on the Symposium of the IAG Subcommission for Europe (EUREF) held in Dubrovnik, May 2001, Torres J. A; Hornik H (eds); Bundesamt für Kartographie und Geodäsie, Frankfurt, Germany, 2001.*

**Greaves M; Cruddace P R; Fane C S;** GPS and Coordinate Transformations in Great Britain, *Civil Engineering Surveyor, GIS / GPS Supplement, September 2002, Vol. 26, No. 7, 2002.*

**Greaves M; Cruddace P R;** A new coordinate Transformation for Great Britain. *GeoConnexions February 2002.*

**Greaves M; Fane C S;** The British EUREF GB 2001 GPS Campaign, *Report on the Symposium of the IAG Subcommission for Europe (EUREF) held in Ponta Delgada, 5 – 8 June 2002, Torres, J. A., Hornik, H. (eds), Bundesamt für Kartographie und Geodäsie, Frankfurt, Germany. Note – not yet published at time of writing. Expected publication date June 2003. Available on line at [http://www.euref-iag.org/book2002/4\\_2.pdf](http://www.euref-iag.org/book2002/4_2.pdf)*

**Hide C; Moore T; Smith M J;** Adaptive Kalman Filtering for Low Cost INS/GPS, *Proc ION GPS 2002, The 15th Technical Meeting of the Satellite Division of the Institute of Navigation, PP 1143-1147, Portland, USA, September 2002.*

**Hill C J; Moore T; Napier M E;** Rapid Mapping with Post Processed Data from Garmin Handheld Receivers, *Proc ION GPS 2002, The 15th Technical Meeting of the Satellite Division of the Institute of Navigation, 9 pages, Portland, USA, September 2002.*

**Hill C J; Moore T; Dumville M;** Carrier Phase Surveying with Garmin Handheld GPS Receivers, *Survey Review, Vol 36, No 280, ISSN 0039-6265, pp 135-141, April 2001.*

**Hill C J; Moore T; Napier M E;** Rapid Mapping with Post-processed Garmin data, *Proc NAV 01, Location and Navigation, pp 13-1 – 13-9, London, November 2001.*

**Hill C J; Moore T;** GPS Surveying without Selective Availability, *Surveying World, Vol 8, Issue 5, ISSN 0927-7900, pp 21 - 22, July / August 2000.*

**Hill C J; Moore T; Ashkenazi V;** Satellite Positioning Methods, *Continental Shelf Limits: The Scientific and Legal Interface, Oxford University Press, ISBN 0-19-511782-4, pp111 - 123, 2000.*

**Hill C J; Moore T; Dumville M;** Gringo: Recording RINEX Data from Handheld GPS Receivers, *Proc GNSS 2000, 4th European Conference on Global Navigation Satellite Systems, pp 12, Edinburgh, May 2000.*

**Hill C J; Moore T; Dumville M;** Carrier Phase Surveying with Garmin Handheld GPS Receivers, *Proc ION GPS 2000, 13th International Technical Meeting of the Satellite Division of the Institute of Navigation, 8 pages, Salt Lake City, USA, 2000.*

**Hill C J; Dumville M; Moore T;** GRINGO: A RINEX Logger for Hand-Held GPS Receivers, *Proc ION-GPS-99, The 12th International Technical Meeting of the Satellite Division of the Institute of Navigation, 8 pages, Nashville, USA, September 1999.*

**Hill C J; Moore T;** Developments in Satellite Navigation Systems, *Surveying World, Vol 7, Issue 4, ISSN 0927-7900, pp 20 - 22, May/June 1999.*

- Hill C J; Moore T; Ashkenazi V;** Satellite Positioning and Navigation, Sovereign Limits Beneath the Oceans: Delimiting the New Continental Shelf, *IHO/IOC, In Print 1999.*
- Hill C J; Moore T; Dumville M D;** Gringo Comes to Town - Improving the Accuracy of Handheld GPS, *Surveying World, Vol 8, Issue 1, ISSN 0927-7900, pp 28 - 30, November / December 1999.*
- Hilton R D; Berry P A M; Featherstone W;** Validation and comparison of digital elevation Models over Australia using Satellite Radar Altimetry, *BGAS meeting, Herstmonceux castle, 2001.*
- Hipkin R G;** Vertical datum defined by  $W_o = U_o$ : theory and practice of a modern height system, *paper presented at the International Gravity & Geoid Commission Meeting, 26-30 August 2002, Thessaloniki. (in press). Also presented at the Weikko A. Heiskanen Symposium in Geodesy, Columbus, Ohio, 1-4 October 2002*
- Hipkin R G;** Earth's gravity spectrum, Kaula's Rule and geoid computation, *paper presented at the Weikko A. Heiskanen Symposium in Geodesy, Columbus, Ohio, 1-4 October 2002b.*
- Hipkin R G;** Ocean-wide gravity network adjustment, *paper presented at 3<sup>rd</sup> Meeting of the International Gravity and Geoid Commission, Thessaloniki, 26-30 August 2002c.*
- Hipkin R G;** Is there a need for a Geodetic Datum 2000? Discussion of a Heiskanen & Moritz proposition, *pp 124 – 127 in Vistas for the New Millenium, IAG Symposium 125, (ed J. Adam & P. Schwarz), Springer-Verlag, Berlin, 2002a.*
- Hipkin R G;** New rationale for local geoid computation from EGM96 covariance modelling, *paper presented at IAG 2001 Scientific Assembly, Budapest. 2-7 September 2001b.*
- Hipkin R G; Haines K; Beggan C; Bingley R M; Woodworth P L; Baker T F; Flather R; Hernandez F;** EDIN2000: A Rationale for Local Geoid Computation and its Realisation, *Proceedings of XXVI General Assembly of the European Geophysical Society, Nice, France, March 2001.*
- Hipkin R G;** The statistics of pink noise on a sphere: application to mantle density anomalies, *Geophysical Journal International, 144, 259-270, 2001a.*
- Hughes C W;** Torques exerted by a shallow fluid on a non-spherical, rotating planet, *Tellus 54A, 56-62, 2002.*
- Hughes C W;** Zonal jets in and near the Coral Sea, seen by satellite altimetry, *Geophys Res. Letters, 29(9) 44-1 - 44-4. DOI 10.1029/2001GL014006, 2002.*
- Hughes C W; Ash E R;** Eddy Forcing of the mean flow in the Southern Ocean, *J. Geophys. Res. (Oceans), 106(C2), 2713-2722, 2001.*
- Hughes C W; Wunsch C; Zlotnicki V;** Satellite Peers through the Oceans from Space, *Eos (Transactions of the American Geophysical Union) 81(7), p68, 2000.*
- Hughes C W; Baker T F;** Oceanography from gravity measurements, A.70. *JSG14/L/11-A3 in, IUGG 99 Birmingham, Abstracts. Week A, Birmingham, Monday 19 July to Saturday 24 July, IUGG 1999.*
- Hughes C W; Carnochan S;** Rossby wave propagation in the Southern Ocean from satellite altimetry, *Advances in Space Research 22(11) 1557-1560, 1999.*
- Husson V; Stevens P; Appleby G M;** System Performance Assessment from NERC Simultaneous Arc Analysis, *Proceedings of 12<sup>th</sup> International Workshop on Laser Ranging, Matera, Italy, November 2000.*
- Iliffe J C; Griffiths W J; Message E L;** Localised Geoid Determination for Engineering Control Surveys, *Survey Review, Vol. 35, No. 275, pp 320 - 328, 2000.*
- ILRS Analysis Working Group;** Collaborating with research groups from some 10 institutes towards an official, regular, ILRS station coordinate and EOP product. *Co-authored a review paper to the 2002 Spring AGU meeting and a 'Call for Participation' paper for distribution to the ILRS analysis community.*
- Jameson P; Walsh D; Brodin G; Cross P; Ziebart M;** High accuracy attitude determination for small satellites using GPS/GNSS signals, *Proceedings of NAVITEC2001 – First ESA Workshop on Satellite Navigation User Equipment Technologies, ESTEC, Noordwijk, The Netherlands, 8pp 2001.*
- Jentzsch G; Baker T F (Eds);** Special section on earth tides and ocean tide loading, *Journal of Physics and Chemistry of the Earth, 25(4): 395-415, 2000.*
- Johannessen J A; Balmino G; Le Provost C; Rummel R; Sabadini R; Sünkel H; Tscherning C C; Visser P; Woodworth P L; Hughes C; LeGrand P; Sneeuw N; Perosanz F; Aguirre-Martinez M; Rebhan H; Drinkwater M;** The European Gravity Field and Steady-State Ocean Circulation Explorer Satellite Mission: Impact in Geophysics, *Reviews of Geophysics - accepted as of 12/12/2002.*

**Johannessen J A; Le Provost C; Drange H; Srokosz M; Woodworth P L; Schlüssel P; Le Grand P; Kerr Y; Wingham D; Rebhan H;** Observing the Ocean from Space: Emerging Capabilities in Europe, pp.198-208 in, *Observing the Oceans in the 21st Century*, C.J. Koblinsky and N.R. Smith (eds), Bureau of Meteorology, Melbourne, Australia. 604pp., 2001.

**Johannessen J; Le Provost C; Drange H; Srokosz M; Woodworth P L; Schlüssel P; Le Grand P; Kerr Y; Wingham D; Rebhan H;** Emerging new Earth Observation capabilities in the context of ocean observing systems for climate, *Proceedings of the conference on The Ocean Observing System for Climate*, St.Raphael, France, 18-22 October, (Solicited papers), 1999.

**Johnson C P D; Berry P A M;** Cross-calibration and validation of altimeter backscatter over natural land targets. *European Geophysical Assembly, Nice France, April 2002.*

**Johnson C P D; Berry P A M;** Validation and correction of space derived Digital Elevation Models using ERS satellite altimeter data. *IAG scientific assembly September 2001.*

**Johnson C P D;** ERS and ENVISAT backscatter monitoring using natural land targets. *BGAS meeting, Herstmonceux castle, 2001.*

**Kadzim M R;** Large scale, high precision GPS network for satellite laser ranging calibration, *MSc Report, Department of Geomatic Engineering, UCL, 2002.*

**Kealy A; Scott-Young S; Leahy F; Cross P;** Intelligent Navigation, Inertial Navigation: Double Assistance for GPS, *GPS World, Vol 13, No 5 (May), pp26-33, 2002.*

**Kealy A; Scott-Young A; Leahy F; Cross P;** Improving the Performance of Satellite Navigation Systems for Land Mobile Applications through the Integration of MEMS Inertial Sensors, *Proceedings of ION-GPS 2001, Salt Lake City, Institute of Navigation pp1394-1402, 2001.*

**Keenan C R;** Stochastic modelling for high fidelity DGPS quality assessment, *PhD thesis University College London, 2001.*

**Kilby G;** Satellite altimeter calibration and precise orbit determination of GFO, *Ph.D. thesis, University of Newcastle upon Tyne, 2001.*

**Lavallée D A;** Tectonic plate motions from global GPS measurements, *Ph.D. thesis, University of Newcastle upon Tyne, 2000.*

**Lythe M B; Vaughan D G; BEDMAP Consortium;** BEDMAP - bed topography of the Antarctic. 1:10,000,000 scale map, *BAS (Misc) 9. Cambridge, British Antarctic Survey, 2000.*

**Mahmud M R;** Precise three-dimensional attitude estimation from independent GPS arrays, *PhD thesis, University College London, 1999.*

**Majumdar A; Ochieng W Y;** Estimation of European Airspace Capacity from a Model of Controller Workload. *Journal of Navigation, 55(3), 381-403, 2002.*

**Majumdar A; Ochieng W Y;** Factors affecting air traffic controller workload: a multivariate analysis based upon simulation modelling of controller workload. *Transportation Research Record, TRR 1788, 58-69, 2002.*

**Majumdar A; Ochieng W Y;** From Our Air is not for sale to Airtrack: The Part Privatisation of the UK's Airspace. *Accepted for publication in the Transport Review Journal, 2002.*

**Mather J S B; Smith M J; Jowett S; Gibson K A H; Moynihan D;** Application of a Photogrammetric Technique to Golf Club Evaluation, *Sports Engineering, Vol 3, No 1, ISSN 1369-7072, pp 37 - 47, 2000.*

**Mathers E L; Woodworth P L;** Departures from the local inverse barometer model observed in altimeter and tide gauge data and in a global barotropic numerical model, *Journal of Geophysical Research, 106(C4), 6957- 6972, 2001.*

**Meng X; Roberts G W; Dodson A H; Cosser E; Noakes C;** Simulation of the Effects of Introducing Pseudolite Data into Bridge Deflection Monitoring Data, *Proc Second Symposium on Geodesy for Geotechnical and Structural Engineering, Berlin, Germany, May 21 - 24, ISBN 3-9501492-1-X, pp 372-381, 2002.*

**Meng X; Roberts G; Dodson A H; Cosser E;** The Use of Pseudolites to Augment GPS Data for Bridge Deflection Measurements. *Proceedings of the 15th International Technical Meeting of the Satellite Division of the Institute of Navigation, Portland, Oregon, USA, pp. 851-862, September 2002.*

**Meng X;** Real-Time Monitoring of Structures using GPS/Accelerometers, Smart Bridge: Concept to Development, *PhD thesis, University of Nottingham, 2002.*

**Meng X; Roberts G W; Dodson A H;** Error Analysis and Accuracy Improvements to GPS/Accelerometer Structural Monitoring, *Proc IAG 2001 Scientific Assembly, Budapest, Hungary, 2-8 September 2001.*

**Meng X;** Vehicle-Borne Highway Geometric Alignment and Facilities Data Dapture by using DGPS and GPS/GIS Integration, *Proc ION-GPS-99, The 12th International Technical Meeting of the Satellite Division of the Institute of Navigation*, 8 pages, Nashville, USA, September 1999.

**Mitchum G T; Cheney R; Fu L-L; Le Provost C; Menard Y; Woodworth P L;** The future of sea surface height observations, pp.120-136 in, *Observing the Oceans in the 21st Century*, C.J. Koblinsky and N.R. Smith (eds), Bureau of Meteorology, Melbourne, Australia. 604pp., 2001.

**Monteiro L N S; Moore T; Hill C J;** Maritime DGPS: Ensuring the Best Availability and Continuity, *ENC GNSS 2000, Proceedings of the European Navigation Conference*, 7 pages, Copenhagen, Denmark, May 2002.

**Moore P; Sterlini P;** Cross-calibration of TOPEX/Poseidon, ERS-2 and GEOSAT altimetry, *Advances in Space Research*, 30(2), 233-239, 2002.

**Moore P; Kilby G;** Orbital inferences for GFO with applications to GEOSAT, *Advances in Space Research*, 30 (2), 393-399, 2002.

**Moore P;** The ERS-2 altimetric bias and gravity field enhancement using dual crossovers between ERS and TOPEX/Poseidon, *Journal of Geodesy*, 75(5-6), 241-254, 2001.

**Moore P and Murphy CM,** Inter-calibrations of multi-satellite altimetric missions, *Advances in Space Research*, 25, 1099-1102, 2000.

**Moore T;** Ionospheric Scintillation Effects on GPS Measurements, *Proc CGSIC ISC European Meeting, Brussels, Belgium*, December 2002.

**Moore T; Aquino M H O; Waugh S; Dodson A H; Hill C J;** Evaluation of the EGNOS Ionospheric Correction Model under Scintillation in Northern Europe, *Proc ION GPS 2002, The 15th Technical Meeting of the Satellite Division of the Institute of Navigation*, 10 Pages, Portland, USA, September 2002.

**Moore T; Aquino M H O; Dodson A H; Waugh S J;** Impact of Ionospheric Scintillation on EGNOS and DGPS Users in Northern Europe, *ENC GNSS 2002, Proceedings of the European Navigation Conference*, 10 pages, 2002.

**Moore T; Bonnor N;** GNSS Vulnerability Tutorials, *Proc NAV 02, International Conference of the Royal Institute of Navigation*, pp 1-88, London, November 2002.

**Moore T; Hill C J; Monteiro L N S;** Maritime DGPS: Ensuring the Best Availability and Continuity, *Journal of Navigation*, Vol 55, No 3, ISSN 0373 4633, pp 485 - 494, September 2002.

**Moore T; Hill C J; Monteiro L N S;** Is DGPS Still a Good Option for Mariners?, *The IALA Bulletin, Magazine of the International Association of Marine Aids to Navigation and Lighthouse Authorities*, 2002/3, ISSN 0379-2811, pp 37 - 43, November 2002.

**Moore T; Hill C J; Monteiro L N S;** Optimising Availability and Continuity of the Portuguese DGPS Network, *Proc ION GPS 2002, The 15th Technical Meeting of the Satellite Division of the Institute of Navigation*, 7 pages, Portland, USA, September 2002.

**Moore T; Roberts G W; Veneboer T; Pattinson M; Close G; Moore R;** New Developments in River Level Monitoring using GPS Heighting, *Proc ION GPS 2002, The 15th Technical Meeting of the Satellite Division of the Institute of Navigation*, 8 pages, Portland, USA, September 2002.

**Moore T;** Directions 2002, *GPS World Showcase*, Vol 12, No 12, p25, December 2001.

**Moore T; Roberts G W; Close G; Moore R;** River Level Monitoring using GPS Heighting, *Proc UN/USA Workshop on the Use and Applications of Global Navigation Satellite Systems*, Vienna, Austria, November 2001.

**Moore T; Dodson A H; Aquino M H O; Waugh S;** Ionospheric Scintillation Monitoring in Northern Europe, *Proc ION GPS 2001, 14th Technical Meeting of the Satellite Division of the Institute of Navigation*, 9 pages, Salt Lake City, USA, September 2001.

**Moore T; Hill C J; Monteiro L N S;** Is DGPS still a Good Option for Mariners?, *Journal of Navigation*, Vol 54, No 3, ISSN 0373-4633, September 2001.

**Moore T; Hill C J; Monteiro L N S;** Is DGPS still a Good Option for Mariners?, *Proc ION GPS 2001, 14th Technical Meeting of the Satellite Division of the Institute of Navigation*, 7 pages, Salt Lake City, USA, September 2001.

**Moore T;** Satellite Navigation Group, *Navigation News*, ISSN 0298 6317, pp 26, May / June 2001.

**Moore T; Ashkenazi V; Dumville M;** Implementation of WGS 84 and GPS for Marine Navigation, *International Hydrographic Review, GITC*, Vol 2, No 1, pp 62-71, 2001.

**Moore T; Hill C J; Monteiro L N S;** The Portuguese DGPS Network, *Proc GNSS 2001, 5th International Conference on Global Navigation Satellite Systems*, Seville, Spain, May 2001.

**Moore T;** Directions 2001, *GPS World Showcase*, Vol 11, No 12, ISSN 1048-5104, 2000.

**Moore T; Close G; Lee C; Moore R;** RiGHt on the Water: River Level Monitoring using GPS Heighting, *Galileo's World*, ISSN 1528-2341, pp 22 - 27, December 2000.

**Moore T; Chen W; Ashkenazi V; Dumville M;** Implementation of WGS84 and GPS for Marine Navigation, *Proc 9th European Meeting of the International Information Sub-Committee, Civil GPS Service Interface Committee*, 7 pages, Monte Carlo, Monaco, December 2000.

**Moore T; Hill C J;** Scientific Applications of Galileo, *Proc NAV 2000*, pp 11-1 - 11-7, London, November 2000.

**Moore T; Chen W; Ashkenazi V; Dumville M;** WGS84 and GNSS for UK Marine Navigation, *Proc ION GPS 2000, 13th International Technical Meeting of the Satellite Division of the Institute of Navigation*, 8 pages, Salt Lake City, USA, September 2000.

**Moore T; Dumville M; Chen W;** Implications of the use of GNSS and WGS84 for Marine Navigation, *Journal of Navigation*, Vol 53, No 3, ISSN 0373-4633, pp 413 - 424, September 2000.

**Moore T; Close G; Moore R;** RiGHt: River Level Monitoring using GPS Heighting, *Proc ION GPS 2000, 13th International Technical Meeting of the Satellite Division of the Institute of Navigation*, 8 pages, Salt Lake City, USA, September 2000.

**Moore T; Hill C J;** The Removal of Selective Availability: The Impact on the Geodetic Community, *The Quarterly Newsletter of the Institute of Navigation*, Vol 10, No 3, pp 4 - 13, Fall 2000.

**Moore T; Zhang K; Close G; Moore R;** Real Time River Level Monitoring using GPS Heighting, *GPS Solutions*, Vol 4, No 2, ISSN 1080-5370, pp 63 - 67, Fall 2000.

**Moore T; Close G; Moore R;** The RiGHt Project: River Level Monitoring using GPS Heighting, *Proc 35th MAFF Conference of River and Coastal Engineers 2000*, pp 11.01.1 - 11.01.4, Keele, July 2000.

**Moore T; Close G; Moore R; Zhang K;** RiGHt: River Level Monitoring using GPS Heighting, *Proc GNSS 2000, 4th European Conference on Global Navigation Satellite Systems*, Edinburgh, 2000.

**Moore T; Chen W; Ashkenazi V; Dumville M;** WGS84 and GNSS for Marine Navigation in the UK, *Proc GNSS 2000, 4th European Conference on Global Navigation Satellite Systems*, 10 pages, Edinburgh, May 2000.

**Moore T;** Contribution to Directions 2000, *GPS World Showcase*, Vol 10, No 12, ISSN 1048-5104, pp 22, December 1999.

**Moore T; Roberts G W; Zhang K F;** Real Time GPS Monitoring of River Levels, *Proc XXII General Assembly of the International Union of Geodesy and Geophysics*, Birmingham, July 1999.

**Moore T; Roberts G W; Zhang K F; Ashkenazi V; Dumville M; Close G; Lee C; Moore R; Tindall I;** A Novel GPS River Level Monitoring System, *Proc GNSS 99, 3rd European Symp on Global Navigation Satellite Systems*, pp 507-512, Genoa, Italy, October 1999.

**Moore T; Roberts G W; Zhang K; Ashkenazi V; Dumville M; Close G; Moore R;** A Real Time GPS River Level Monitoring System, *Proc ION-GPS-99, The 12th International Technical Meeting of the Satellite Division of the Institute of Navigation*, 8 pages, Nashville, USA, September 1999.

**Moore T; Roberts G W ;** Carrier Phase GPS Navigation to the North Pole, *Journal of Navigation*, Vol 52, No 1, ISSN 0373-4633, pp 80 - 89, January 1999.

**Nash J; Latham N; Oakley T; Bingley R M; Waugh S J;** Examination of Integrated Water Vapour Fields Associated with Organised Thunderstorm Outbreaks over the UK in July 2001. *Proceedings of the Workshop of European COST Action 716*, Potsdam, Germany, January 2002.

**Noomen R; Appleby G M; Shelus P J;** The Role of ILRS Within IGGOS, *AGU, Washington*, June 2002.

**Novak R W;** Local-Tie Survey of Satellite Laser Ranging Facility, Herstmonceux, *MSc Report, Department of Geomatic Engineering, UCL*, 2002.

**Ochieng W Y; Sauer K; Sheridan K; Cross P A; Lannelongue S; Ammour N; Petit K;** Integrity mechanisms for GPS satellites within the Galileo architecture, *Journal of Geospatial Engineering*, 4(2), 95-110, 2002.

**Ochieng W Y; Sauer K;** Urban Road Transport Navigation: Performance of GPS after Selective Availability. *The International Journal of Transportation Research - Part C on Emerging Technologies*, 10(3), 171-187, 2002.

**Ochieng W Y; Sheridan K; Han X; Cross P A; Lannelongue S; Ammour N; Petit K;** An Assessment of the RAIM Performance of a Combined Galileo/GPS Navigation System using the Marginally Detectable Errors (MDE) Algorithm. *The GPS Solutions Journal*, 5(3), 42-51, 2002.

**Ochieng W Y; Sheridan K; Han X; Cross P A; Lannelongue S; Ammour N; Petit K;** Integrity Performance Models for a Combined Galileo/GPS Navigation System. *Journal of Geospatial Engineering*, 3(1), 21-32, 2001.



**Ochieng W Y; Sauer K; Cross P A; Sheridan K; Iliffe J; Lannelongue S; Ammour N; Petit K;** Potential performance levels of a combined Galileo/GPS navigation system, *Journal of Navigation*, 54 (2), 186-197, 2001.

**Ochieng W Y; Chen W;** High Level Architectural and Programmatic Issues of the Galileo System. *Journal of Geospatial Engineering*, 2(2), 16-20, 2000.

**Ochieng W Y; Shardlow P J; Johnston G;** Advanced Transport Telematics Positioning Requirements: An Assessment of GPS Performance in Greater London. *Journal of Navigation*, 52(3), 342-355, 1999.

**Ogundipe O; Roberts G W; Dodson A H;** Construction Plant Control and Guidance - The Development Continues. *Civil Engineering Surveyor GIS Supplement*, pp 4 - 9, Autumn 2002, ISSN 0266-139X, 2002.

**Otsubo T; Appleby G M;** Recovery of target response function for centre-of-mass corrections of spherical satellites, *Proceedings of 13<sup>th</sup> International Workshop on Laser Ranging, Washington DC, October 2002* and accepted, *JGR*.

**Otsubo T; Appleby G M; Gibbs P;** GLONASS laser ranging accuracy with satellite signature effect, *Surveys in Geophysics* 22 (6): 509-516 2001.

**Otsubo T; Gibbs P; Walters D; Wood R;** Determination of Lageos Spin Motion from Photometric Observation, *Proceedings of 12<sup>th</sup> International Workshop on Laser Ranging, Matera, Italy, November 2000*.

**Otsubo T; Appleby G M; Gibbs P;** Systematic Range Bias Related to GLONASS Reflector Array, *Proceedings of 12<sup>th</sup> International Workshop on Laser Ranging, Matera, Italy, November 2000*.

**Otsubo T; Appleby G M;** Spectral analysis of SLR full-rate data for timer problem detection, *Laser Radar Ranging and Atmospheric Lidar Techniques II, SPIE Vol. 3865, 33-37, 1999*.

**Page D;** Real-time kinematic and fast static GPS measurements of ground deformation on Mount Etna, Sicily, *Ph.D. thesis, University of Newcastle upon Tyne, 2000*.

**Park D W G;** Using Linear Features for Absolute and Exterior Orientation, *PhD thesis, University of Nottingham, 1999*.

**Pattinson M; Dodson A H; Moore T;** Tropospheric Delay Estimation from a Moving Platform, *Proc ION GPS 2002, The 15th Technical Meeting of the Satellite Division of the Institute of Navigation, 9 pages, Portland, USA, September 2002*.

**Pattinson M;** Estimation of Tropospheric Delay from a Moving GPS Receiver, *PhD thesis, University of Nottingham, 2002*.

**Penna N T; Baker T F;** Ocean tide loading considerations for GPS processing around Australia, *Geomatics Research Australasia*, 77, 1-26, 2002.

**Penna N T; Bingley R M; Dodson A H;** Single receiver Heighting using the Active Stations of the GPS Network of Great Britain, *Survey Review, Vol 36, No 283, pp 340-350, 2002*.

**Penna N T; Dodson A H; Chen W;** Assessment of EGNOS tropospheric correction model, *Journal of Navigation, Vol 54, No 1, ISSN 0373-4633, pp 37-55, 2001*.

**Penna N T; Dodson A H; Chen W;** Assessment of EGNOS Tropospheric Correction Model, *Proc GNSS 2000, 4th European Conference on Global Satellite Navigation Satellite Systems, 18 pages, Edinburgh, May 2000*.

**Pinnock R A; Berry P A M;** Global scale hydrology from satellite altimeter missions: current status and future prospects. *European Geophysical Assembly, Nice France, April 2002*.

**Pinnock R A; Berry P A M;** Potential for global scale hydrological monitoring using ERS altimetry. *IAG scientific assembly September 2001*.

**Pinnock R A; Berry P A M;** Hydrological Applications Of Land Altimetry. *European Geophysical Assembly, Nice France, April 2001*.

**Pinnock R A; Berry P A M;** ERS RA waveform shapes and implications for the ENVISAT mode switching algorithm. *BGAS meeting, Herstmonceux castle, 2001*.

**Pinnock R A; Berry P A M;** A study of regional hydrological monitoring using ERS altimetry. *BGAS meeting, Herstmonceux castle, 2001*.

**Pinnock R A; Berry P A M;** Environmental signals in satellite altimeter data. *BGAS meeting, Met. Office training college Reading, United Kingdom, 1999*.

**Price W F;** Monitoring of Glulam Structures by Theodolite Intersection, *PhD thesis, University of Nottingham, 2002*.

**Rainbow J; Clarke P;** Optimising navigational precision from potential GNSS constellations. *The Hydrographic Journal* 103, 3-9, 2002.

**Raper S C B; Woodworth P L et al;** Global changes in the volume and mass of the ocean, pp. 9-80 in *Sea Level Change and Coastal Processes: Implications for Europe* (eds. D. Smith, S. Raper, S. Zerbini, and A. Sanchez-Arcilla), Luxembourg: Office for Official Publications of the European Communities. 247pp. 2000.

**Rintoul S R; Hughes C W; Olbers D J;** The Antarctic Circumpolar Current System, *Chapter 4.6, pp 271-302 in Ocean Circulation and Climate, Observing and Modelling the Global Ocean, G. Siedler, J. Church, and J. Gould (Eds.), Academic Press, 715 pp., 2001.*

**Roberts G; Ogundipe O; Dodson A H;** Construction Plant Control Using RTK GPS: Case Study. *Proceedings of the 15th International Technical Meeting of the Satellite Division of the Institute of Navigation, Portland, Oregon, USA, pp. 1813-1818, September 2002.*

**Roberts G; Cosser E; Meng X; Dodson A H;** Creating Virtual L2 GPS Data Using a Combination of L1 Receivers and L1/L2 Reference Receivers. *Proceedings of the 15th International Technical Meeting of the Satellite Division of the Institute of Navigation, Portland, Oregon, USA, September 2002, pp 1408-1413, 2002.*

**Roberts G W; Meng X; Dodson A H;** Using Adaptive Filtering to Detect Multipath and Cycle Slips in GPS/Accelerometer Bridge Deflection Monitoring Data, *Proc XXII International Congress of the FIG, TS6.2 Engineering Surveys for Construction Works and Structural Engineering II, Washington DC, USA, April 19 - 26 2002.*

**Roberts G W; Evans A J; Dodson A H; Denby B; Cooper S; Hollands R;** The Use of Augmented Reality, GPS and INS for Subsurface Data Visualisation, *Proc XXII International Congress of the FIG, TS5.13 Integration of Techniques, Washington DC, USA, April 19 - 26 2002.*

**Roberts G W; Strange K; Waller M;** The Detection of Abandoned Mineshafts using GPS and Capacitively Coupled Resistivity Imaging, *Proc XXII International Congress of the FIG, JS27 Engineering Survey Databases and Facility Management Systems, Washington DC, USA, April 19 - 26 2002.*

**Roberts G W; Ogundipe O; Dodson A H;** Construction Plant Control Using RTK GPS, *Proc XXII International Congress of the FIG, TS6.2 Engineering Surveys for Construction Works and Structural Engineering II, Washington DC, USA, April 19 - 26 2002.*

**Roberts G W; Evans A J; Dodson A H; Denby B; Cooper S; Hollands R;** Look Beneath the Surface with Augmented Reality, *GPS World, ISSN 1048-5104, February 2002.*

**Roberts G W; Meng X; Dodson A H; Cosser E;** Geodetic Signal Diagnosis and its Applications to Structural Deformation, *Proc Second Symposium on Geodesy for Geotechnical and Structural Engineering, Berlin, Germany, May 21 - 24 2002, ISBN 3-9501492-1-X, pp 111-122, 2002.*

**Roberts G W; Meng X; Dodson A H; Cosser E;** Multipath Mitigation for Bridge Deformation Monitoring, *the International Journal of GPS, <http://www.cpgps.org/journal/journal.html>, ISSN 1446-3156, 2002.*

**Roberts G W; Meng X; Dodson A H;** Data Processing and Multipath Mitigation Approaches for GPS/Accelerometer Based Hybrid Structural Deflection Monitoring System, *Proc ION-GPS-01, The 14th International Technical Meeting of the Satellite Division of the Institute of Navigation, Salt Lake City, USA, September 2001.*

**Roberts G W; Evans A J; Dodson A H; Denby B; Hollands R; Cooper S;** Integrating GPS, INS and Augmented Reality for sub-surface visualisation, *Proc ION-GPS-01, The 14th International Technical Meeting of the Satellite Division of the Institute of Navigation, Salt Lake City, USA, September 2001.*

**Roberts G W; Meng X; Dodson A H;** The Use of Kinematic GPS and Triaxial Accelerometers to Monitor the Deflections of Large Bridges, *Proc Deformation Measurements and Analysis, 10th International Symposium on Deformation Measurements; International Federation of Surveyors (FIG); Commission 6 - Engineering Surveys; Working Group 6.1, Orange, USA, March 2001.*

**Roberts G W; Meng X; Dodson A H;** Structural Dynamic and Deflection Monitoring using Integrated GPS and Tri-axial Accelerometers, *Proc ION GPS 2000, 13th International Technical Meeting of the Satellite Division of the Institute of Navigation, 8 pages, Salt Lake City, USA, September 2000.*

**Roberts G W; Dodson A H; Ashkenazi V;** Experimental Plant Guidance and Control by Kinematic GPS, *New Civil Engineer International, pp 30 -36, June 2000.*

**Roberts G W; Dodson A H; Ashkenazi V;** Experimental Plant Guidance and Control by Kinematic GPS, *Proc of the ICE (Civil Engineering), No 138, ISSN 0965 089 X, pp 19 - 25, 2000.*

**Roberts G W; Dodson A H; Brown C J; Karuna R; Evans E;** Monitoring the Height Deflections of the Humber Bridge by GPS, GLONASS and Finite Element Modelling, *IAG Symposia Geodesy Beyond 2000, Springer-Verlag, Vol 121, ISBN 3-540-67002-5, pp 355 - 360, Berlin, 2000.*

**Roberts G W; Ashkenazi V; Dodson A H;** Monitoring the height deflections of the Humber Bridge by GPS and GLONASS, *Proc XXII General Assembly of the International Union of Geodesy and Geophysics, Birmingham, July 1999.*

**Roberts G W; Dodson A H;** Using RTK GPS to control construction plant, *Civil Engineering Surveyor, ISSN 0266-139X, pp 24-26, November 1999.*

**Roberts G W; Dodson A H; Ashkenazi V;** Twist and Deflect: Monitoring motion of the Humber Bridge, *GPS World, Vol 10, No 10, ISSN 1048-5104, pp 24-34, October 1999.*

- Roberts G W; Dodson A H; Ashkenazi V; Brown C J; Karuna R; Evans E;** The Use of Kinematic GPS and Finite Element Modelling for the Deformation Measurements of the Humber Bridge, *Proc GNSS 99, 3rd European Symp on Global Navigation Satellite Systems*, pp 230-235, Genoa, Italy, October 1999.
- Roberts G W; Dodson A H; Ashkenazi V; Brown C J; Karuna R; Evans E;** Comparison of GPS Measurements and Finite Element Modelling for Deformation Measurements of the Humber Bridge, *Proc ION-GPS-99, The 12th International Technical Meeting of the Satellite Division of the Institute of Navigation*, 8 pages, Nashville, USA, September 1999.
- Roberts G W; Ashkenazi V; Dodson A H;** Height Control of Construction Plant by GPS and GLONASS, *Proc FIG 5 symposium on the Importance of Heights in Surveying, LMV-Rapport 1999:3, ISSN 0280-5731*, pp 195-204, Gävle, Sweden, March 1999.
- Roberts G W; Dodson A H; Ashkenazi V;** GPS-Aided Autonomous Construction Plant Control and Guidance, *Automation in Construction, Elsevier, Vol 8, ISSN 0926-5805*, 1999.
- Rummel R; Balmino G; Johannessen J A; Visser P; Woodworth P L;** Dedicated gravity field missions - principles and aims, *Journal of Geodynamics*, 33, 3-20, 2002.
- Russell D C;** DEM Creation for Application in Precision Agriculture, *PhD thesis, University of Nottingham*, 2001.
- Russell D C; Smith M J; Dodson A H; Steven M D; Stafford J V;** DEM Creation using GPS for use in Precision Farming, *Proc GNSS 2000, 4th European Conference on Global Navigation Satellite Systems*, 11 pages, Edinburgh, May 2000.
- Russell D C; Smith M J; Dodson A H; Steven M D; Stafford J V;** Assessment of GLA DGPS System for Land Operations within the UK, *Proc GNSS 2000, 4th European Conference on Global Navigation Satellite Systems*, Edinburgh, May 2000.
- Russell D C; Dodson A H; Smith M J; Steven M D; Stafford J V;** Elevation Modelling using GPS for use in Precision Agriculture, *Proc GIS Brasil 99, V Latin American Congress for users of Geoprocessing*, Salvador, Brazil, July 1999.
- Russell D C; Dodson A H; Smith M J; Steven M D; Stafford J V;** Elevation Modelling using GPS and Digital Photogrammetry for use in Precision Agriculture, *Proc Precision Agriculture '99, 2nd European Conference on Precision Agriculture*, Odense, Denmark, July 1999.
- Sanli D U; Blewitt G;** Geocentric sea level trend using GPS and >100-year tide gauge record on a postglacial rebound nodal line. *J. Geophys. Res.*, 106(B1), 713719, 2001.
- Sanli D U;** GPS strategies for tide gauge monitoring with assessment of sea level analysis models, *Ph.D. thesis, University of Newcastle upon Tyne*, 1999.
- Sauer K; Ochieng W Y;** Integrated use of GPS and EGNOS Carrier Phase Observations for High Precision Kinematic Positioning. *Journal of Geospatial Engineering*, 4(1), 59-67, 2002.
- Schwahn W; Baker T F; Falk R; Jeffries G; Lothhammer A; Richter B; Wilmes H; Wolf P;** Long-term increase of gravity at the Medicina Station (Northern Italy) confirmed by absolute and superconducting gravimetric time series, 145-168 in, *Proceedings of the Workshop: High precision gravity measurements with applications to geodynamics and Second GGP Workshop, March 24th to 26th, 1999, Ducarme, B. and J. Barthélemy, Eds. Munsbach Castle, Grand-Duchy of Luxemburg, March 24th to 26th, 1999, Conseil de l'Europe, Cahiers du Centre Européen de Géodynamique et de Séismologie. Volume 17. 2000.*
- Sheridan K F;** GPS based position and attitude determination for airborne remote sensing, *PhD thesis University College London*, 2001.
- Sheridan K F; Ochieng W Y; Saur K; Cross P A; J Iliffe J; Lannelongue S; Ammour N; Petit K;** Performance Potential of a Combined Galileo/GPS Navigation System, *Proceedings of GNSS2001, Seville, Spain, 8-11 May, 2001.*
- Sheridan K; Paul C; Mohd R M;** GPS based attitude determination for airborne remote sensing. In *Geodesy Beyond 2000: the Challenges of the First Decade. International Association of Geodesy Symposia, Springer, Vol 121, pp337-342, 2000.*
- Sheridan K F; Cross P A;** A robust and reliable GPS-based system for the direct georeferencing of airborne line scanning sensors, *Proceedings of GNSS 2000, Edinburgh, 2000.*
- Sherwood R S; Wood R; Otsubo T;** LAGEOS-2 spin rate and orientation, *Proceedings of 13<sup>th</sup> International Workshop on Laser Ranging, Washington DC, October 2002.*
- Shukla R;** Referencing of Airborne Imagery using Direct Measurement of Position and Attitude, *PhD thesis, University of Nottingham*, 2001.
- Shukla R; Smith M J;** Geo-Referencing CASI Imagery using Direct Measurement of Position and Attitude, *International Archives of Photogrammetry and Remote Sensing, ISPRS Congress 2000, Vol XXXIII, Part Book 2, pp 502 - 509, Amsterdam, The Netherlands, July 2000.*

**Shum C K; Zhao C Y; Tseng H Z; Woodworth P L;** Twentieth century sea level change in the Pacific basin, pp.83-88 in, *Proceedings of the Pacific Islands Conference on Climate Change, Climate Variability and Sea Level Rise, 3-7 April, 2000, National Auditorium, Rarotonga, Cook Islands.* (eds. M.Grzechnik and J.Chittleborough). Adelaide: National Tidal Facility Australia. 88pp, 2001.

**Smith A J E; Ambrosius B A C; Wakker K F; Woodworth P L; Vassie J M;** Ocean tides from harmonic and response analysis on TOPEX/POSEIDON altimetry. (Paper presented at the 31st Scientific Assembly of COSPAR, Birmingham UK, 14-21 July 1996). *Advances in Space Research, 22, 1541-1548, 1999.*

**Smith M J; Shukla R;** Providing control for geo-referencing of airborne imagery, *The 5th International Airborne Remote Sensing Conference and Exhibition, San Francisco, USA 17-20 September 2001.*

**Smith M J; Priestnall G; Asal F G;** Evaluating Digital surface models from LIDAR and digital photogrammetry, *The 5th International Airborne Remote Sensing Conference and Exhibition, San Francisco, USA 17-20 September 2001.*

**Smith M J; Park D W G;** Developments in Exterior Orientation of Aerial Photographs, *Canadian Journal of Remote Sensing, Vol 26, No. 5, ISSN 0703-8992, pp 475 - 479, October 2000.*

**Smith M J; Asal F; Priestnall G;** Combining LiDAR and Photogrammetry for Urban and Rural Landscape Studies, *International Archives of Photogrammetry and Remote Sensing, ISPRS Congress 2000, Vol XXXIII, Prt Book 3A, Amsterdam, The Netherlands, July 2000.*

**Smith M J; Park D W G;** Absolute and Exterior Orientation using Linear Features, *International Archives of Photogrammetry and Remote Sensing, ISPRS Congress 2000, Vol XXXIII, Prt Book 3B , pp 850 - 857, Amsterdam, The Netherlands, July 2000.*

**Smith M J; Park D W G;** A Case for Indirect Image Orientation using Linear Features, ISPRS Commission III Workshop Theory and Algorithms WG III/1, *Integrated Sensor Calibration and Orientation, pp 53-62, Barcelona, Spain, November 1999.*

**Smith M J; Park D W G;** Towards a New Approach for Absolute and Exterior Orientation, *Photogrammetric Record, Vol XVI, No 94, pp 617-623, October 1999.*

**Smith M J;** Review of Digital Landscape Model for Europe (DLME) by E Blau et al, *Photogrammetric Record, Vol XVI, No 94, pp 729-730, October 1999.*

**Smith M J;** Navigation Satellites and Airborne Images for Earth Observation, *Journal of the Nottingham Society of Engineers, Vol 7, No 2, pp 12-13, 1999.*

**Smith M J; Park D W G;** Developments in Exterior Orientation of Aerial Photographs, *Proc Fourth International Airborne Remote Sensing Conference and Exhibition, pp 501-508, Ottawa, Canada, June 1999.*

**Smith M J; Park D W G;** Advances in Exterior Orientation, *ASPRS Annual Convention and Exhibition, pp 877-883, Portland, USA, April 1999.*

**Sterlini P E;** Global sea surface height changes from satellite altimetry, *Ph.D. thesis, University of Newcastle upon Tyne, 2001.*

**Stow R J; Boyle J; Wright P; Bingley R M; Hobbs P; Crilly M;** London Clay Shrink/Swell Deformation Maps by SAR Interferometry, with Atmospheric Phase Correction derived from Rain Radar and Validation by a Regional GPS Network, *Proceedings of XXVI General Assembly of the European Geophysical Society, Nice, France, March 2001.*

**Strange K; Roberts G; Waller M;** The Detection of Abandoned Mineshafts Using GPS and Capacitively Coupled Resistivity Imaging. *Proceedings of the 15th International Technical Meeting of the Satellite Division of the Institute of Navigation, Portland, Oregon, USA, September 2002, pp 1423-1428, 2002.*

**Swann J W;** Advantages and Problems of Combining GPS and GLONASS, *PhD thesis, University of Nottingham, 1999.*

**Teferle F N; Bingley R M; Dodson A H; Baker T F;** Application of the Dual-GPS Concept to Monitoring Vertical Land Movements at Tide Gauges, *Journal of Physics and Chemistry of the Earth, Vol 27/32-34, pp 1401-1406, 2002.*

**Teferle F N; Bingley R M; Dodson A H; Penna N T; Baker T F;** Using GPS to Separate Crustal Movements and Sea Level Changes at Tide Gauges in the UK, Vertical Reference Systems, Ed. Drewes, H, *International Association of Geodesy Symposia, Vol 124, pp 264-269, ISBN 3-540-43011-3, Springer Verlag, 2002.*

**Teferle F N; Bingley R M; Dodson A H; Baker T F; Williams S D P;** Evaluation of the Dual-CGPS Concept to Monitoring Vertical Land Movements at Tide Gauges, *Proceedings of the International Association of Geodesy 2001 Scientific Assembly, Budapest, Hungary, September 2001.*

**Teferle F N; Bingley R M; Dodson A H; Baker T F;** Application of the Dual-CGPS Concept to Monitoring Vertical Land Movements at Tide Gauges, *Proceedings of XXVI General Assembly of the European Geophysical Society, Nice, France, March 2001.*

- Teferle F N**; Continuous GPS Measurements at UK Tide Gauge Sites; 1997 - 2000, *Proc ION GPS 2000, 13th International Technical Meeting of the Satellite Division of the Institute of Navigation*, 8 pages, Salt Lake City, USA, September 2000.
- Tsimplis M N; Baker T F**; Sea level drop in the Mediterranean Sea: an indicator of deep water salinity and temperature changes? *Geophysical Research Letters*, 27(12): 1731-1734, 2000.
- Tsou Y-L**; Ionospheric modelling for high precision GPS real time kinetic surveying over long distances, *PhD thesis University College London*, 2000.
- Turner J F**; Precise orbit determination and gravity field recovery of CHAMP, *Ph.D. thesis, University of Newcastle upon Tyne*, 2002.
- Visser P N A M; Rummel R; Balmino G; Sunkel H; Johannessen J; Aguirre M; Woodworth P L; Le Provost C; Tscherning C C; Sabadini R**; The European Earth Explorer Mission GOCE: impact for the geosciences, pp.95-107 in, *Ice Sheets, Sea Level and the Dynamic Earth. Geodynamics Series 29, American Geophysical Union*, eds. J. Mitrovica and L.L.A. Vermeersen, 2002.
- Vitushkin L; Becker M; Jiang Z; Francis O; van Dam T M; Faller J; Chartier J M; Amalvict M; Bonvalot S; Debeglia N; Desogus S; Diament M; Dupont F; Falk R; Gabalda G; Gagnon C G L; Gattacceca T; Germak A; Hinderer J; Jamet O; Jeffries G; Kaker R; Kopaev A; Liard J; Lindau A; Longuevergne L; Luck B; Maderal E N; Makinen J; Meurers B; Mizushima S; Mrlina J; Newell D; Origlia C; Pujol E R; Reinhold A; Richard P; Robinson I A; Ruess D; Thies S; van Camp M; van Ruymbeke M; Compagni, M F D; Williams S**; Results of the Sixth International Comparison of Absolute Gravimeters, *ICAG-2001, Metrologia*, 39 (5), 407-424, 2002.
- Vrhovski D; Moore T; Bennett L D**; GPS Vulnerability in Urban Positioning for , *Proc NAV 02 GNSS Vulnerability, International Conference of the Royal Institute of Navigation*, 8 pages, London, November 2002.
- Wadge G; Webley P W; James I N; Bingley R; Dodson A; Waugh S; Veneboer T; Puglisi G; Mattia M; Baker D F; Edwards S C; Edwards S J; Clarke P J**;; Atmospheric models, GPS and InSAR measurements of the tropospheric water vapour field over Mount Etna. *Geophys. Res. Lett.*, 29(19), 1905, doi:10.1029/2002GL015159, 2002
- Webley P J; Bingley R M; Dodson A H; Wadge G; Waugh S J; James I N**; Atmospheric Water Vapour Correction to InSAR Surface Motion Measurements on Mountains: Results from a Dense GPS Network on Mount Etna, *Physics and Chemistry of the Earth, Vol 27, Issue 4-5*, pp 363-370, 2002.
- Webley P J; Bingley R M; Dodson A H; Wadge G; Waugh S J; James I N**; The Use of GPS Integrated Water Vapour (IWV) in Meteorological Modelling and Synthetic Aperture Radar Interferometry (InSAR), *Proceedings of XXVI General Assembly of the European Geophysical Society, Nice, France, March 2001*.
- Williams S D P; Baker T F; Jeffries G**; Absolute gravity measurements at UK tide gauges, *Geophysical Research Letters*, 28 (12), 2317-2320, 2001.
- Wilson I; Cruddace P R; Pennington G; Greaves M; Fane C S; Euler H J; Keenan R; Levine M**; Establishing an Network RTK Solution in Great Britain; From Schema to Solution. *Proceedings of ION GPS 2002 conference, Portland, Oregon*, 2002.
- Wolf R; Rossbach U; Werner W [IfEN GmbH]; Lannelongue S; Michelin J [Alcatel Space Industries]; Thomas P [Logica UK Ltd.]**; EGNOS CPF Check Set - Stand Alone Performance, *Proceedings of ION GPS 2002, Portland, Oregon*.
- Wood R; Gurtner W**; Berne/Herstmonceux Timebias Service, *Proceedings of 13<sup>th</sup> International Workshop on Laser Ranging, Washington DC, October 2002*.
- Woodworth P L; Le Provost C; Rickards L J; Mitchum G T; Merrifield M**;; A review of sea-level research from tide gauges during the World Ocean Circulation Experiment, *Oceanography and Marine Biology: An Annual Review*, 40, 1-35, 2002
- Woodworth P L; Blackman D L**; Changes in extreme high waters at Liverpool since 1768, *International Journal of Climatology*, 22, 697-714, 2002
- Woodworth P L**; (ed.) Manual on sea-level measurement and interpretation. Volume III: Reappraisals and recommendations as of the year 2000, *Intergovernmental Oceanographic Commission, Manuals & Guides, No 14*, 47pp. & appendices, 2002.
- Woodworth P L; Gregory J M**; Benefits of GRACE and GOCE to sea level studies, *Space Science Reviews (in press)* 2002.
- Woodworth P L**; Report by the Permanent Service for Mean Sea Level (PSMSL) for the period 1999-2001 to the IAG Scientific Assembly, Budapest, September 2001, *Proceedings of the Assembly of the International Association of Geodesy (O.B. Andersen, ed.)*, IAG Travaux volume 32, IAG central bureau, Copenhagen, 2001. (To be published as a book and cdrom, available also at <http://www.gfy.ku.dk/~iag/Travaux/contents.htm>), 2001.
- Woodworth P L**; Monitoring and predicting long term sea level changes, *Periodicum Biologorum Vol. 102, Supplement 1*, 665-672, 2000.

**Woodworth P L; Tsimplis M N; Flather R A; Shennan I;** A review of the trends observed in British Isles mean sea level data measured by tide gauges, *Geophysical Journal International*, 136, 651-670, 1999.

**Woodworth P L;** High waters at Liverpool since 1768: the UK's longest sea level record, *Geophysical Research Letters*, 26 (11), 1589-1592, 1999.

**Woodworth P L;** A study of changes in high water levels and tides at Liverpool during the last two hundred and thirty years with some historical background, *Proudman Oceanographic Laboratory Report No.56, 62pp. & figures*, 1999.

**Wright T J; Parsons B E; Jackson J A; Haynes M; Fielding E J; England P C; Clarke P J;** Source parameters of the 1 October 1995 Dinar (Turkey) earthquake from SAR interferometry and seismic body wave modelling. *Earth Planet. Sci. Lett.*, 172, 23-37, 1999.

**Zerbini S; Baker T F; Cazenave A; Cocard A M; Kaas E; Kahle H G; Lozano I; Marson I; Plag H P; Romagnoli H; von Storch H;** Regional and local sea level variations, 81-133 in, *Sea level change and coastal processes: implications for Europe*. Smith D; Raper S B; Zerbini S; Sanchez-Arcilla A (Eds). Luxembourg, 2000.

**Zhang K F; Dodson A H; Chen W;** Factors Affecting FFT Gravimetric Geoid Determination Precision, *Phys. Chem. Earth (A)*, Vol 25, No 1, ISSN 1464-1895, pp 31-37, 2000.

**Zhang K F; Dodson A H;** Precision Evaluation of the Terrain Corrections and Geoid Determination through FFT Technique, presented to the 24th General Assembly of the European Geophysical Society, The Hague, The Netherlands, April 1999.

**Zhang K F; Dodson A H; Chen W;** Factors Affecting FFT Gravimetric Geoid Determination Precision, *Physics and Chemistry of the Earth, In Print* 1999.

**Ziebart M; Cross P; Adhya S;** Modelling Photon Pressure - the Key to High Precision GPS Satellite Orbits, *GPS World*, Vol. 13, No.1, p43-50, 2002.

**Ziebart M; Dare P;** Analytical Radiation Pressure Modelling for GLONASS using a Pixel Array, *Journal of Geodesy*, 75: 587-599, 2001.